# Sialin (S-17): sc-50975



The Power to Question

# **BACKGROUND**

Sialin, also designated sodium/sialic acid cotransporter or membrane glycoprotein HP59, belongs to the major facilitator superfamily and the sodium/anion cotransporter family of proteins. Sialin acts as a solute translocator for anionic substances; in lysosomes it is a free sialic acid transporter. Sialin is a multi-pass membrane protein that localizes to the lysosome and is primarily detected in fetal lung and small intestine. It is also expressed in adult placenta, kidney and pancreas, and may be detected in colon, breast and ovary tumor endothelial cells as well. Defects in the SLC17A5 gene, which encodes the Sialin protein, can cause several disorders including Salla disease, an autosomal recessive sialic acid storage disease, as well as infantile sialic acid storage disorder (ISSD), a severe form of sialic acid storage disease in which affected newborns exhibit coarse features, visceromegaly (an enlargement of the internal organs) and a failure to thrive after birth.

# **REFERENCES**

- Biancheri, R., et al. 2005. Homo-zygosity for the p.K136E mutation in the SLC17A5 gene as cause of an Italian severe Salla disease. Neurogenetics 6: 195-199.
- Allen, S., et al. 2005. Novel sialic acid transporter of *Haemophilus influenzae*. Infect. Immun. 73: 5291-5300.
- 3. Post, D.M., et al. 2005. Identification of a novel sialic acid transporter in *Haemophilus ducreyi*. Infect. Immun. 73: 6727-6735.
- Yarovaya, N., et al. 2005. Sialin, an anion transporter defective in sialic acid storage diseases, shows highly variable expression in adult mouse brain, and is developmentally regulated. Neurobiol. Dis. 19: 351-365.
- Wreden, C.C., et al. 2005. Varied mechanisms underlie the free sialic acid storage disorders. J. Biol. Chem. 280: 1408-1416.
- Morse, R.P., et al. 2006. Novel form of intermediate Salla disease: clinical and neuroimaging features. J. Child Neurol. 20: 814-816.

# CHROMOSOMAL LOCATION

Genetic locus: SLC17A5 (human) mapping to 6q13; Slc17a5 (mouse) mapping to 9 E1.

# SOURCE

Sialin (S-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Sialin of human origin.

## **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50975 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **APPLICATIONS**

Sialin (S-17) is recommended for detection of Sialin of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Sialin siRNA (h): sc-61545, Sialin siRNA (m): sc-61546, Sialin shRNA Plasmid (h): sc-61545-SH, Sialin shRNA Plasmid (m): sc-61546-SH, Sialin shRNA (h) Lentiviral Particles: sc-61545-V and Sialin shRNA (m) Lentiviral Particles: sc-61546-V.


Molecular Weight of Sialin: 55 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204.

# **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **DATA**



Sialin (S-17): sc-50975. Western blot analysis of Sialin expression in Jurkat whole cell lysate.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com