

# Glyoxalase II (C-17): sc-51091

## BACKGROUND

The glyoxal pathway plays a role in the detoxification of glucose degradation products (GDP). Glyoxalase I and Glyoxalase II (also designated hydroxyacyl glutathione hydrolase or HAGH) are members of the Glyoxalase family. The Glyoxalase II enzyme is a thiolesterase that catalyzes the hydrolysis of S-D-lactoyl-glutathione to form reduced glutathione and D-lactic acid. It exists only as a monomer and binds two zinc ions per subunit. Glyoxalase II contains 260 amino acids. It is detected in the mitochondria and cytosol of mammals. Both Glyoxalase I and Glyoxalase II are detected at a higher activity level in breast cancer tissues than with matched unaffected tissues. This suggests that Glyoxalase inhibitor drugs may be effective in the treatment of cancer.

## REFERENCES

- Ridderström, M., et al. 1996. Molecular cloning, heterologous expression and characterization of human Glyoxalase II. *J. Biol. Chem.* 271: 319-323.
- Cameron, A.D., et al. 1999. Crystal structure of human Glyoxalase II and its complex with a glutathione thiolester substrate analogue. *Structure* 7: 1067-1078.
- Rulli, A., et al. 2001. Expression of Glyoxalase I and II in normal and breast cancer tissues. *Breast Cancer Res. Treat.* 66: 67-72.
- Cordell, P.A., et al. 2004. The human hydroxyacylglutathione hydrolase (HAGH) gene encodes both cytosolic and mitochondrial forms of Glyoxalase II. *J. Biol. Chem.* 279: 28653-28661.
- Krömer, S.A., et al. 2005. Identification of Glyoxalase I as a protein marker in a mouse model of extremes in trait anxiety. *J. Neurosci.* 25: 4375-4384.
- Yadav, S.K., et al. 2005. Methylglyoxal levels in plants under salinity stress are dependent on Glyoxalase I and glutathione. *Biochem. Biophys. Res. Commun.* 337: 61-67.
- Ariza, A., et al. 2006. Specificity of the trypanothione-dependent *Leishmania major* Glyoxalase I: structure and biochemical comparison with the human enzyme. *Mol. Microbiol.* 59: 1239-1248.

## CHROMOSOMAL LOCATION

Genetic locus: HAGH (human) mapping to 16p13.3; Hagh (mouse) mapping to 17 A3.3.

## SOURCE

Glyoxalase II (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Glyoxalase II of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51091 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Glyoxalase II (C-17) is recommended for detection of Glyoxalase II of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Glyoxalase II (C-17) is also recommended for detection of Glyoxalase II in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Glyoxalase II siRNA (h): sc-60705, Glyoxalase II siRNA (m): sc-60706, Glyoxalase II shRNA Plasmid (h): sc-60705-SH, Glyoxalase II shRNA Plasmid (m): sc-60706-SH, Glyoxalase II shRNA (h) Lentiviral Particles: sc-60705-V and Glyoxalase II shRNA (m) Lentiviral Particles: sc-60706-V.


Molecular Weight of Glyoxalase II: 29 kDa.

Positive Controls: Glyoxalase II (m2): 293T Lysate: sc-120531 or mouse liver extract: sc-2256.

## DATA



Glyoxalase II (C-17): sc-51091. Western blot analysis of Glyoxalase II expression in non-transfected: sc-117752 (A) and mouse Glyoxalase II transfected: sc-120531 (B) 293T whole cell lysates.



Glyoxalase II (C-17): sc-51091. Immunofluorescence staining of formalin-fixed HepG2 cells showing cytoplasmic localization.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **Glyoxalase II (F-9): sc-166781** or **Glyoxalase II (F-12): sc-365025**, our highly recommended monoclonal alternatives to Glyoxalase II (C-17).