

MTMR5 (I-13): sc-51135

BACKGROUND

Myotubularin and the myotubularin-related proteins (MTMR1-9) belong to a highly conserved family of eukaryotic phosphatases. They are protein tyrosine phosphatases that utilize inositol phospholipids, rather than phosphoproteins, as substrates. MTMR family members hydrolyze both phosphatidylinositol 3-phosphate (PtdIns3P) and PtdIns(3,5)P₂. MTMR2 interacts with MTMR5, an inactive family member that increases the enzymatic activity of MTMR2 and dictates its subcellular localization. Mutations in MTMR2 cause autosomal recessive Charcot-Marie-Tooth type 4B1 (CMT4B1), which is characterized by reduced nerve conduction velocities, focally folded myelin sheaths and demyelination. MTMR3 and MTMR4 can either interact with each other or self-associate. MTMR6 regulates the activity of the calcium-activated potassium channel 3.1. MTMR9 regulates the activity of MTMR7 and MTMR8.

REFERENCES

1. Laporte, J., et al. 1997. Mutations in the MTM1 gene implicated in X-linked myotubular myopathy. *Hum. Mol. Genet.* 6: 1505-1511.
2. Blondeau, F., et al. 2000. Myotubularin, a phosphatase deficient in myotubular myopathy, acts on phosphatidylinositol 3-kinase and phosphatidylinositol 3-phosphate pathway. *Hum. Mol. Genet.* 9: 2223-2229.
3. Kim, S.A., et al. 2003. Regulation of myotubularin-related MTMR2 phosphatidylinositol phosphatase by MTMR5, a catalytically inactive phosphatase. *Proc. Natl. Acad. Sci. USA* 100: 4492-4497.
4. Mochizuki, Y. and Majerus, P.W. 2003. Characterization of myotubularin-related protein 7 and its binding partner, myotubularin-related protein 9. *Proc. Natl. Acad. Sci. USA* 100: 9768-9773.
5. Srivastava, S., et al. 2005. Phosphatidylinositol 3-phosphate indirectly activates KCa3.1 via 14 amino acids in the carboxy terminus of KCa3.1. *Mol. Biol. Cell* 17: 146-154.
6. Lorenzo, O., et al. 2006. Systematic analysis of myotubularins: heteromeric interactions, subcellular localisation and endosomerelated functions. *J. Cell Sci.* 119: 2953-2959.

CHROMOSOMAL LOCATION

Genetic locus: SBF1 (human) mapping to 22q13.33; Sbf1 (mouse) mapping to 15 E3.

SOURCE

MTMR5 (I-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MTMR5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51135 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

MTMR5 (I-13) is recommended for detection of MTMR5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MTMR5 siRNA (h): sc-61092, MTMR5 siRNA (m): sc-63318, MTMR5 shRNA Plasmid (h): sc-61092-SH, MTMR5 shRNA Plasmid (m): sc-63318-SH, MTMR5 shRNA (h) Lentiviral Particles: sc-61092-V and MTMR5 shRNA (m) Lentiviral Particles: sc-63318-V.

Molecular Weight of MTMR5: 208 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **MTMR5 (B-9): sc-393488** or **MTMR5 (B-3): sc-393565**, our highly recommended monoclonal alternatives to MTMR5 (I-13).