

TRESK (H-17): sc-51235

BACKGROUND

Potassium channels play an important role in cell excitability and plasticity. The pore loop domain, a highly conserved region common to all potassium channels, is involved in determining potassium ion selectivity. The family of potassium channels possessing two-pore loop domains consists of both inward and outwardly rectifying channels and includes THIK-1, THIK-2, TRESK, TALK-1 and TALK-2. Members of this family, which are all characterized by four transmembrane domains, may function to help influence the resting membrane potential of cells. TWIK-related spinal cord K⁺ (TRESK) is the most sensitive volatile anesthetic-activated channel in the family and may function to mediate the effects of inhaled anesthetics in the central nervous system in a manner that is sensitive to immunosuppressive drugs. TRESK is activated by the calcium signal from calcineurin, a calcium/calmodulin-dependent phosphatase, and is highly sensitive to zinc.

REFERENCES

- Czirják, G., et al. 2004. The two-pore domain K⁺ channel, TRESK, is activated by the cytoplasmic calcium signal through calcineurin. *J. Biol. Chem.* 279: 18550-18558.
- Kang, D., et al. 2004. Functional expression of TRESK-2, a new member of the tandem-pore K⁺ channel family. *J. Biol. Chem.* 279: 28063-28070.
- Liu, C., et al. 2004. Potent activation of concentrations of volatile anesthetics. *Anesth. Analg.* 99: 1715-1722.
- Kang, D., et al. 2005. Thermosensitivity of the two-pore domain K⁺ channels TREK-2 and TRAAK. *J. Physiol.* 564: 103-116.
- Keshavaprasad, B., et al. 2005. Species-specific differences in response to anesthetics and other modulators by the K2P channel TRESK. *Anesth. Analg.* 101: 1042-1049.
- Brosnan, R., et al. 2006. Chirality in anesthesia II: stereoselective modulation of ion channel function by secondary alcohol enantiomers. *Anesth. Analg.* 103: 86-91.

CHROMOSOMAL LOCATION

Genetic locus: KCNK18 (human) mapping to 10q25.3.

SOURCE

TRESK (H-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TRESK of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51235 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TRESK (H-17) is recommended for detection of TRESK of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TRESK siRNA (h): sc-61709, TRESK shRNA Plasmid (h): sc-61709-SH and TRESK shRNA (h) Lentiviral Particles: sc-61709-V.


Molecular Weight of TRESK: 43 kDa.

Positive Controls: TRESK (h): 293T Lysate: sc-158006, HeLa whole cell lysate: sc-2200 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TRESK (H-17): sc-51235. Western blot analysis of TRESK expression in non-transfected 293T: sc-117752 (A), human TRESK transfected 293T: sc-158006 (B), Hep G2 (C) and HeLa (D) whole cell lysates and mouse brain tissue extract (E).



TRESK (H-17): sc-51235. Immunofluorescence staining of formalin-fixed HepG2 cells showing membrane localization.

SELECT PRODUCT CITATIONS

- Sánchez-Miguel, D.S., et al. 2013. TRESK potassium channel in human T lymphoblasts. *Biochem. Biophys. Res. Commun.* 434: 273-279.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.