

Twinfilin-1 (G-16): sc-51242

BACKGROUND

Twinfilin is a highly conserved Actin monomer-binding protein that regulates cytoskeletal dynamics in organisms from yeast to mammals. Twinfilin is composed of two ADF-homology domains; it coordinates filament severing and monomer sequestering at sites of rapid Actin turnover, thus preventing assembly of the monomer into filaments. Twinfilin-1 is the mammalian homolog and is expressed in embryos and in most adult non-muscle cell types. Twinfilin-1 binds ADP-G-Actin and efficiently halts Actin filament assembly by inhibiting the nucleotide exchange on Actin monomers and directly interacting with the capping protein. Phosphatidylinositol (4,5)-bisphosphate inhibits the activity of Twinfilin-1 and Rac1 and Cdc42, two small GTPases, induce the redistribution of Twinfilin-1 to membrane ruffles and cell-cell contacts, respectively.

REFERENCES

- Goode, B.L., Drubin, D.G. and Lappalainen, P. 1998. Regulation of the cortical Actin cytoskeleton in budding yeast by Twinfilin, a ubiquitous Actin monomer-sequestering protein. *J. Cell Biol.* 142: 723-733.
- Vartiainen, M., Ojala, P.J., Auvinen, P., Peränen, J. and Lappalainen, P. 2000. Mouse A6/Twinfilin is an Actin monomer-binding protein that localizes to the regions of rapid Actin dynamics. *Mol. Cell. Biol.* 20: 1772-1783.
- Palmgren, S., Ojala, P.J., Wear, M.A., Cooper, J.A. and Lappalainen, P. 2001. Inter activity and localization of yeast Twinfilin. *J. Cell Biol.* 155: 251-260.
- Wahlström, G., Vartiainen, M., Yamamoto, L., Mattila, P.K., Lappalainen, P. and Heino, T.I. 2001. Twinfilin is required for Actin-dependent developmental processes in *Drosophila*. *J. Cell Biol.* 155: 787-796.
- Palmgren, S., Vartiainen, M. and Lappalainen, P. 2002. Twinfilin, a molecular mailman for Actin monomers. *J. Cell. Sci.* 115 (Pt 5): 881-886.
- Vartiainen, M.K., Sarkkinen, E.M., Matilainen, T., Salminen, M. and Lappalainen, P. 2003. Mammals have two Twinfilin isoforms whose subcellular localizations and tissue distributions are differentially regulated. *J. Biol. Chem.* 278: 34347-34355.
- Falck, S., Paavilainen, V.O., Wear, M.A., Grossmann, J.G., Cooper, J.A. and Lappalainen, P. 2004. Biological role and structural mechanism of Twinfilin-capping protein interaction. *EMBO J.* 23: 3010-3019.
- Rush, J., Moritz, A., Lee, K.A., Guo, A., Goss, V.L., Spek, E.J., Zhang, H., Zha, X.M., Polakiewicz, R.D. and Comb, M.J. 2005. Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. *Nat. Biotechnol.* 23: 94-101.
- Moseley, J.B., Okada, K., Balcer, H.I., Kovar, D.R., Pollard, T.D. and Goode, B.L. 2006. Twinfilin is an Actin-filament-severing protein and promotes rapid turnover of Actin structures *in vivo*. *J. Cell. Sci.* 119 (Pt 8): 1547-1557.

CHROMOSOMAL LOCATION

Genetic locus: PTK9 (human) mapping to 12q12; Ptk9 (mouse) mapping to 15 F1.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

Twinfilin-1 (G-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Twinfilin-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51242 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Twinfilin-1 (G-16) is recommended for detection of Twinfilin-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Twinfilin-1 siRNA (h): sc-61738.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.