



WIT-1 (N-12): sc-51247

BACKGROUND

Wilms' tumor is a neoplasm of the kidneys that usually occurs in children and is characterized by the presence of abortive tubules and glomeruli surrounded by a spindle cell stroma. The 11p13 Wilms tumor locus consists of two coordinately regulated transcripts, WT1 and WIT-1, which are mutated in Wilms' tumors. Wilms' tumor upstream neighbor 1 (WIT-1) is encoded by an intronless gene upstream of the Wilms' tumor 1 (WT1) gene, which is important for nephrogenesis and gonadal growth. The WT1 gene is bi-directionally transcribed from the same promoter region as WIT-1, which may function as an antisense regulator of WT1. WIT-1 and WT1 have the same temporal and cell-restricted expression pattern, although the expression of WIT-1 is less abundant. Methylation of the WIT-1 gene is implicated in hematologic malignancy of chemoresistant acute myeloid leukemia. Single nucleotide polymorphisms (SNPs) in the WIT-1 gene are significantly associated with focal segmental glomerulosclerosis.

REFERENCES

- Huang, A., Campbell, C.E., Bonetta, L., McAndrews-Hill, M.S., Chilton-MacNeill, S., Coppes, M.J., Law, D.J., Feinberg, A.P., Yeger, H. and Williams, B.R. 1990. Tissue, developmental, and tumor-specific expression of divergent transcripts in Wilms' tumor. *Science* 250: 991-994.
- Coppes, M.J., Bonetta, L., Huang, A., Hoban, P., Chilton-MacNeill, S., Campbell, C.E., Weksberg, R., Yeger, H., Reeve, A.E. and Williams, B.R. 1993. Loss of heterozygosity mapping in Wilms' tumor indicates the involvement of three distinct regions and a limited role for nondisjunction or mitotic recombination. *Genes Chromosomes Cancer* 5: 326-334.
- Gessler, M. and Bruns, G.A. 1993. Sequence of the WT1 upstream region including the WIT-1 gene. *Genomics* 17: 499-501.
- Eccles, M.R., Grubb, G., Ogawa, O., Szeto, J. and Reeve, A.E. 1994. Cloning of novel Wilms' tumor gene (WT1) cDNAs; evidence for antisense transcription of WT1. *Oncogene* 9: 2059-2063.
- Hewitt, J.A., Kessler, P.M., Campbell, C.E. and Williams, B.R. 1996. Tissue-specific regulation of the WT1 locus. *Med. Pediatr. Oncol.* 27: 456-461.
- Plass, C., Yu, F., Yu, L., Strout, M.P., El-Rifai, W., Elonen, E., Knuutila, S., Marcucci, G., Young, D.C., Held, W.A., Bloomfield, C.D. and Caligiuri, M.A. 1999. Restriction landmark genome scanning for aberrant methylation in primary refractory and relapsed acute myeloid leukemia; involvement of the WIT-1 gene. *Oncogene* 18: 3159-3165.
- Orloff, M.S., Iyengar, S.K., Winkler, C.A., Goddard, K.A., Dart, R.A., Ahuja, T.S., Mokrzycki, M., Briggs, W.A., Korbet, S.M., Kimmel, P.L., Simon, E.E., Trachtman, H., Vlahov, D., Michel, D.M., Berns, J.S., Smith, M.C., Schelling, J.R., Sedor, J.R. and Kopp, J.B. 2005. Variants in the Wilms' tumor gene are associated with focal segmental glomerulosclerosis in the African American population. *Physiol. Genomics* 21: 212-221.

CHROMOSOMAL LOCATION

Genetic locus: WIT1 (human) mapping to 11p13.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

WIT-1 (N-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of WIT-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51247 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

WIT-1 (N-12) is recommended for detection of WIT-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for WIT-1 siRNA (h): sc-61802.

Molecular Weight of WIT-1: 10 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.