

# GRWD1 (D-1): sc-514042

## BACKGROUND

GRWD1 (glutamate-rich WD repeat-containing protein 1), also known as WDR28 or KIAA1942, is a 446 amino acid protein that contains five WD repeats. Localizing to the nucleus, GRWD1 is a member of the 50S and 80S preribosomal complexes and may play a role in ribosome biogenesis. The gene encoding GRWD1 maps to human chromosome 19q13.33 and mouse chromosome 7 B4. Consisting of around 63 million bases with over 1,400 genes, human chromosome 19 makes up over 2% of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG families, and Fc $\alpha$  receptors. Peutz-Jeghers syndrome, spinocerebellar ataxia type 6, the stroke disorder CADASIL, hypercholesterolemia and Insulin-dependent diabetes have been linked to chromosome 19.

## REFERENCES

1. Teglund, S., et al. 1994. The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family. *Genomics* 23: 669-684.
2. Wang, L., et al. 2000. C-CAM1, a candidate tumor suppressor gene, is abnormally expressed in primary lung cancers. *Clin. Cancer Res.* 6: 2988-2993.
3. Trowsdale, J., et al. 2001. The genomic context of natural killer receptor extended gene families. *Immunol. Rev.* 181: 20-38.
4. Le Meur, N., et al. 2004. Complete germline deletion of the STK11 gene in a family with Peutz-Jeghers syndrome. *Eur. J. Hum. Genet.* 12: 415-418.

## CHROMOSOMAL LOCATION

Genetic locus: GRWD1 (human) mapping to 19q13.33; Grwd1 (mouse) mapping to 7 B4.

## SOURCE

GRWD1 (D-1) is a mouse monoclonal antibody raised against amino acids 139-318 mapping within an internal region of GRWD1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG $\gamma$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

GRWD1 (D-1) is available conjugated to agarose (sc-514042 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514042 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514042 PE), fluorescein (sc-514042 FITC), Alexa Fluor<sup>®</sup> 488 (sc-514042 AF488), Alexa Fluor<sup>®</sup> 546 (sc-514042 AF546), Alexa Fluor<sup>®</sup> 594 (sc-514042 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-514042 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-514042 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-514042 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

GRWD1 (D-1) is recommended for detection of GRWD1 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GRWD1 siRNA (h): sc-97614, GRWD1 siRNA (m): sc-145787, GRWD1 shRNA Plasmid (h): sc-97614-SH, GRWD1 shRNA Plasmid (m): sc-145787-SH, GRWD1 shRNA (h) Lentiviral Particles: sc-97614-V and GRWD1 shRNA (m) Lentiviral Particles: sc-145787-V.


Molecular Weight of GRWD1: 49 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, HeLa whole cell lysate: sc-2200 or K-562 whole cell lysate: sc-2203.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## DATA



GRWD1 (D-1): sc-514042. Western blot analysis of GRWD1 expression in HeLa (A), K-562 (B), RT-4 (C) and NIH/3T3 (D) whole cell lysates and HeLa nuclear extract (E).



GRWD1 (D-1): sc-514042. Immunofluorescence staining of formalin-fixed SW480 cells showing nuclear speckle and cytoplasmic localization.

## STORAGE

Store at 4 $^{\circ}$  C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.