

SP-lyase (V-15): sc-51432

BACKGROUND

Sphingosine-1-phosphate lyase (SP-lyase) is a member of the group II decarboxylase family that is involved in lipid metabolism. SP-lyase has a variety of functions contributing to normal development, including maintenance of the reproductive system, stress responses, tissue integrity and cell survival. Located in the membrane of the endoplasmic reticulum, SP-lyase is responsible for the irreversible degradation of sphingosine-1-phosphate (S1P). S1P is a lipid important in cell proliferation and migration and, once cleaved by SP-lyase, is degraded into fatty acids and phosphoethanolamine. Through its ability to regulate S1P expression, SP-lyase may play a role in stress-induced apoptosis and is thought to exhibit tumor suppressor activity by silencing S1P activity. Current research suggests that SP-lyase may be a useful target for cancer therapy drugs, as increasing its expression during tumorigenesis may help to regulate cell proliferation.

REFERENCES

- Zhou, J. and Saba, J.D. 1998. Identification of the first mammalian sphingosine phosphate lyase gene and its functional expression in yeast. *Biochem. Biophys. Res. Commun.* 242: 502-507.
- Van Veldhoven, P.P., et al. 2000. Human sphingosine-1-phosphate lyase: cDNA cloning, functional expression studies and mapping to chromosome 10q22. *Biochim. Biophys. Acta* 1487: 128-134.
- Ikeda, M., et al. 2004. Sphingosine-1-phosphate lyase SPL is an endoplasmic reticulum-resident, integral membrane protein with the pyridoxal 5'-phosphate binding domain exposed to the cytosol. *Biochem. Biophys. Res. Commun.* 325: 338-343.
- Oskouian, B., et al. 2005. Regulation of sphingosine-1-phosphate lyase gene expression by members of the GATA family of transcription factors. *J. Biol. Chem.* 280: 18403-18410.
- Min, J., et al. 2005. Sphingosine-1-phosphate lyase regulates sensitivity of human cells to select chemotherapy drugs in a p38-dependent manner. *Mol. Cancer Res.* 3: 287-296.
- Kariya, Y., et al. 2005. Products by the sphingosine kinase/sphingosine 1-phosphate (S1P) lyase pathway but not S1P stimulate mitogenesis. *Genes Cells* 10: 605-615.
- Schwab, S.R., et al. 2005. Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients. *Science* 309: 1735-1739.
- Bassi, R., et al. 2006. Sphingosine-1-phosphate is released by cerebellar astrocytes in response to bFGF and induces astrocyte proliferation through G protein-coupled receptors. *Glia* 53: 621-630.
- Bandhuvula, P. and Saba, J.D. 2007. Sphingosine-1-phosphate lyase in immunity and cancer: silencing the siren. *Trends Mol. Med.* 13: 210-217.

CHROMOSOMAL LOCATION

Genetic locus: SGPL1 (human) mapping to 10q22.1; Sgpl1 (mouse) mapping to 10 B4.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

SP-lyase (V-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of SP-lyase of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51432 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SP-lyase (V-15) is recommended for detection of SP-lyase of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SP-lyase (V-15) is also recommended for detection of SP-lyase in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SP-lyase siRNA (h): sc-72291, SP-lyase siRNA (m): sc-72292, SP-lyase shRNA Plasmid (h): sc-72291-SH, SP-lyase shRNA Plasmid (m): sc-72292-SH, SP-lyase shRNA (h) Lentiviral Particles: sc-72291-V and SP-lyase shRNA (m) Lentiviral Particles: sc-72292-V.

Molecular Weight of SP-lyase: 63 kDa.

Positive Controls: HUV-EC-C whole cell lysate: sc-364180.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.