


# ACOT9 (G-6): sc-514330



The Power to Question

## BACKGROUND

Acyl-CoA thioesterases (ACOTs) are a group of enzymes that catalyze the hydrolysis of acyl-CoA to form coenzyme A (CoA) and a free fatty acid. Through their catalytic activity, ACOTs are able to regulate the level of fatty acids and acyl-CoAs within the cell. ACOT9 (acyl-CoA thioesterase 9), also known as ACATE2, MT-ACT48 (mitochondrial acyl-CoA thioesterase of 48 kDa) or CGI-16, is a 406 amino acid member of the acyl-CoA hydrolase protein family. ACOT9 contains a C-terminal 80 amino acid domain that is conserved from mouse to human, suggesting that the C-terminus may confer the catalytic activity of ACOT9. The gene encoding ACOT9 is located on chromosome X and the expressed ACOT9 protein is localized to the mitochondrion.

## REFERENCES

- Murphy, D.J., et al. 1984. Solubilization, purification and kinetic properties of three membrane-bound long-chain acyl-coenzyme-A thioesterases from microsomes of photosynthetic tissue. *Eur. J. Biochem.* 142: 43-48.
- Alexson, S.E., et al. 1993. Isolation and characterization of microsomal acyl-CoA thioesterase. A member of the rat liver microsomal carboxyl-esterase multi-gene family. *Eur. J. Biochem.* 214: 719-727.
- Wilcke, M. and Alexson, S.E. 1994. Characterization of acyl-CoA thioesterase activity in isolated rat liver peroxisomes. Partial purification and characterization of a long-chain acyl-CoA thioesterase. *Eur. J. Biochem.* 222: 803-811.
- Poupon, V., et al. 1999. Molecular cloning and characterization of MT-ACT48, a novel mitochondrial acyl-CoA thioesterase. *J. Biol. Chem.* 274: 19188-19194.
- Lai, C.H., et al. 2000. Identification of novel human genes evolutionarily conserved in *Caenorhabditis elegans* by comparative proteomics. *Genome Res.* 10: 703-713.
- Hunt, M.C., et al. 2005. A revised nomenclature for mammalian acyl-CoA thioesterases/hydrolases. *J. Lipid Res.* 46: 2029-2032.

## CHROMOSOMAL LOCATION

Genetic locus: ACOT9 (human) mapping to Xp22.11; Acot9 (mouse) mapping to X F3.

## SOURCE

ACOT9 (G-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 11-29 near the N-terminus of ACOT9 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-514330 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

ACOT9 (G-6) is recommended for detection of ACOT9 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ACOT9 siRNA (h): sc-91052, ACOT9 siRNA (m): sc-140816, ACOT9 shRNA Plasmid (h): sc-91052-SH, ACOT9 shRNA Plasmid (m): sc-140816-SH, ACOT9 shRNA (h) Lentiviral Particles: sc-91052-V and ACOT9 shRNA (m) Lentiviral Particles: sc-140816-V.


Molecular Weight of ACOT9: 48 kDa.

Positive Controls: U-87 MG cell lysate: sc-2411, MCF7 whole cell lysate: sc-2206 or HEK293 whole cell lysate: sc-45136.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



ACOT9 (G-6): sc-514330. Western blot analysis of ACOT9 expression in U-87 MG (A), MCF7 (B), HEK293 (C) and MDA-MB-435S (D) whole cell lysates.



ACOT9 (G-6): sc-514330. Western blot analysis of ACOT9 expression in Sol8 (A) and A-10 (B) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.