

CD40 (G-12): sc-514493

BACKGROUND

Resting B cells can be activated and clonally expanded into antibody-producing cells in response to a combination of cell contact and soluble signals provided by primed helper T (Th) cells. While cytokines IL-4 and IL-13 alone are inadequate for B cell activation, contact with Th cells seems to be sufficient for delivery of proliferative signals. A receptor ligand pair central to the transmission of this signal is CD40, expressed on the surface of B cells, together with CD40L, expressed on activated T cells. In the presence of such stimulus, IL-4 and IL-13 are capable of triggering immunoglobulin class switching and secretion of IgE. B cells are sensitive to these cytokines only subsequent to CD40/CD40L-driven DNA synthesis. A downstream mediator of the CD40 signaling pathway, designated CRAF, is a member of an expanding family of proteins that contain a conserved cysteine- and histidine-rich RING finger motif. Other members of the family include TRAF1 and TRAF2. The latter proteins have been shown to regulate TNF-R2 as well as CD40 signaling through activation of the NF κ B family of transcription factors.

REFERENCES

1. Kehry, M.R. and Hodgkin, P.D. 1994. B cell activation by helper T cell membranes. *Crit. Rev. Immunol.* 14: 221-238.
2. Hu, H.M., et al. 1994. A novel RING finger protein interacts with the cytoplasmic domain of CD40. *J. Biol. Chem.* 269: 30069-30072.
3. Rothe, M., et al. 1994. A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* 78: 681-682.

CHROMOSOMAL LOCATION

Genetic locus: CD40 (human) mapping to 20q13.12; Cd40 (mouse) mapping to 2 H3.

SOURCE

CD40 (G-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 242-259 within a C-terminal cytoplasmic domain of CD40 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD40 (G-12) is available conjugated to agarose (sc-514493 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514493 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514493 PE), fluorescein (sc-514493 FITC), Alexa Fluor® 488 (sc-514493 AF488), Alexa Fluor® 546 (sc-514493 AF546), Alexa Fluor® 594 (sc-514493 AF594) or Alexa Fluor® 647 (sc-514493 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-514493 AF680) or Alexa Fluor® 790 (sc-514493 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-514493 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

CD40 (G-12) is recommended for detection of precursor and mature CD40 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CD40 siRNA (h): sc-29250, CD40 siRNA (m): sc-29998, CD40 shRNA Plasmid (h): sc-29250-SH, CD40 shRNA Plasmid (m): sc-29998-SH, CD40 shRNA (h) Lentiviral Particles: sc-29250-V and CD40 shRNA (m) Lentiviral Particles: sc-29998-V.


Molecular Weight of CD40: 43 kDa.

Positive Controls: Raji whole cell lysate: sc-364236, GA-10 whole cell lysate: sc-364230 or BJAB whole cell lysate: sc-2207.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CD40 (G-12): sc-514493. Western blot analysis of CD40 expression in BJAB (A), GA-10 (B) and Raji (C) whole cell lysates.



CD40 (G-12): sc-514493. Western blot analysis of CD40 expression in Ramos whole cell lysate.

SELECT PRODUCT CITATIONS

1. Zhang, L., et al. 2021. Intravenous transplantation of olfactory ensheathing cells reduces neuroinflammation after spinal cord injury via interleukin-1 receptor antagonist. *Theranostics* 11: 1147-1161.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.