

C4BP β (E-1): sc-514553

BACKGROUND

The complement component proteins C3, C4 and C5 are potent anaphylatoxins that are released during classical complement activation, a system of ligand-surface protein interactions that aid in the elimination of pathogens. These proteins belong to the α 2-Macroglobulin family, but retain distinctive features including an anaphylatoxin domain and a netrin (NTR) domain. They are also expressed as single-chain precursors, which are cleaved into α , β and γ subunits that are linked by disulfide bonds. Complement C4 is an essential component for the activation of the complement pathway, which acts through the receptor CR1 (CD35). Complement C4 is predominately expressed in liver and its precursor contains C4a anaphylatoxin and C4b. The full length C4 protein is cleaved into an α chain, a β chain and a γ chain. C4 exists as two functionally distinct isotypes, C4A and C4B, which react preferentially with amino groups and hydroxyl groups, respectively. Excessive complement activation by C4 is negatively regulated by C4BP (C4 binding protein), a fluid-phase complement inhibitor that protects against complement-induced cell apoptosis. The C4BP complex contains α and β chains which act together to accelerate inactivation of C4, thereby controlling the classical pathway of complement activation.

REFERENCES

1. Scharfstein, J., et al. 1978. Human C4-binding protein. I. Isolation and characterization. *J. Exp. Med.* 148: 207-222.
2. Chung, L.P., et al. 1985. Molecular cloning and characterization of the cDNA coding for C4b-binding protein, a regulatory protein of the classical pathway of the human complement system. *Biochem. J.* 230: 133-141.
3. Blom, A.M., et al. 2000. Positively charged amino acids at the interface between α -chain CCP1 and CCP2 of C4BP are required for regulation of the classical C3-convertase. *Mol. Immunol.* 37: 445-453.

CHROMOSOMAL LOCATION

Genetic locus: C4BPB (human) mapping to 1q32.2.

SOURCE

C4BP β (E-1) is a mouse monoclonal antibody raised against amino acids 82-252 mapping at the C-terminus of C4BP β of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

C4BP β (E-1) is available conjugated to agarose (sc-514553 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514553 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514553 PE), fluorescein (sc-514553 FITC), Alexa Fluor[®] 488 (sc-514553 AF488), Alexa Fluor[®] 546 (sc-514553 AF546), Alexa Fluor[®] 594 (sc-514553 AF594) or Alexa Fluor[®] 647 (sc-514553 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-514553 AF680) or Alexa Fluor[®] 790 (sc-514553 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

C4BP β (E-1) is recommended for detection of C4BP β of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for C4BP β siRNA (h): sc-42741, C4BP β shRNA Plasmid (h): sc-42741-SH and C4BP β shRNA (h) Lentiviral Particles: sc-42741-V.


Molecular Weight of C4BP β : 52 kDa.

Positive Controls: human plasma extract: sc-364374, HeLa whole cell lysate: sc-2200 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



C4BP β (E-1): sc-514553. Western blot analysis of C4BP β expression in Hep G2 (A) and HeLa (B) whole cell lysates.



C4BP β (E-1) HRP: sc-514553 HRP. Direct western blot analysis of C4BP β in human plasma. Cruz Marker[™] Molecular Weight Standards detected with Cruz Marker MW Tag-HRP: sc-516732.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.