

APPBP2 (K-20): sc-51460

BACKGROUND

APPBP2 (β -Amyloid precursor protein-binding protein2), also known as protein interacting with APP tail 1 (PAT1) or ARA67, is a hydrophilic, microtubule binding protein that functions in the trafficking of β -Amyloid precursor protein. It is expressed in a variety of cell types and localizes to the cytoplasm. APPBP2 shares homology with Kinesin light chain. It consists of a coiled coil domain, PKC phosphorylation sites, four imperfect C-terminal tandem repeats, eight tetratricopeptide repeats and N- and C-terminal globular structures. APPBP2 recognizes and binds to the basolateral sorting sequence (BaSS) present in the cytoplasmic domain of β -Amyloid precursor protein. In addition, APPBP2 interacts with the androgen receptor and suppresses androgen signaling.

REFERENCES

- Nagase, T., Seki, N., Ishikawa, K., Ohira, M., Kawarabayasi, Y., Ohara, O., Tanaka, A., Kotani, H., Miyajima, N. and Nomura, N. 1997. Prediction of the coding sequences of unidentified human genes. VI. The coding sequences of 80 new genes (KIAA0201-KIAA0280) deduced by analysis of cDNA clones from cell line KG-1 and brain. *DNA Res.* 3: 321-329, 341-354.
- Monni, O., Barlund, M., Mousses, S., Kononen, J., Sauter, G., Heiskanen, M., Paavola, P., Avela, K., Chen, Y., Bittner, M.L. and Kallioniemi, A. 2001. Comprehensive copy number and gene expression profiling of the 17q23 amplicon in human breast cancer. *Proc. Natl. Acad. Sci. USA* 98: 5711-5716.
- Hirasawa, A., Saito-Ohara, F., Inoue, J., Aoki, D., Susumu, N., Yokoyama, T., Nozawa, S., Inazawa, J. and Imoto, I. 2003. Association of 17q21-q24 gain in ovarian clear cell adenocarcinomas with poor prognosis and identification of PPM1D and APPBP2 as likely amplification targets. *Clin. Cancer Res.* 9: 1995-2004.
- Zhang, Y., Yang, Y., Yeh, S. and Chang, C. 2004. ARA67/PAT1 functions as a repressor to suppress androgen receptor transactivation. *Mol. Cell. Biol.* 24: 1044-1057.
- Miyauchi, S., Abbot, E.L., Zhuang, L., Subramanian, R., Ganapathy, V. and Thwaites, D.T. 2005. Isolation and function of the amino acid transporter PAT1 (slc36a1) from rabbit and discrimination between transport via PAT1 and system IMINO in renal brush-border membrane vesicles. *Mol. Membr. Biol.* 22: 549-559.
- Hsu, C.L., Chen, Y.L., Ting, H.J., Lin, W.J., Yang, Z., Zhang, Y., Wang, L., Wu, C.T., Chang, H.C., Yeh, S., Pimplikar, S.W. and Chang, C. 2005. Androgen receptor (AR) NH₂- and COOH-terminal interactions result in the differential influences on the AR-mediated transactivation and cell growth. *Mol. Endocrinol.* 19: 350-361.

CHROMOSOMAL LOCATION

Genetic locus: APPBP2 (human) mapping to 17q23.2; Appbp2 (mouse) mapping to 11 C.

SOURCE

APPBP2 (K-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of APPBP2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-51460 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

APPBP2 (K-20) is recommended for detection of APPBP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

APPBP2 (K-20) is also recommended for detection of APPBP2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for APPBP2 siRNA (h): sc-106762, APPBP2 siRNA (m): sc-141177, APPBP2 shRNA Plasmid (h): sc-106762-SH, APPBP2 shRNA Plasmid (m): sc-141177-SH, APPBP2 shRNA (h) Lentiviral Particles: sc-106762-V and APPBP2 shRNA (m) Lentiviral Particles: sc-141177-V.

Molecular Weight (predicted) of APPBP2: 67 kDa.

Molecular Weight (observed) of APPBP2: 63 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209 or human fetal brain tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfaction
Guaranteed

Try **APPBP2 (4-RE24): sc-134266**, our highly recommended monoclonal alternative to APPBP2 (K-20).