

CREB3L1 (F-8): sc-514635

BACKGROUND

CREB3L1 (cAMP-responsive element-binding protein 3-like protein 1), also designated OASIS (old astrocyte specifically induced substance), is a 519 amino acid transcription factor that activates unfolded protein response target genes during endoplasmic reticulum (ER) stress. CREB3L1 may be specifically involved in the ER-stress response in astrocytes of the central nervous system. CREB3L1 increases inducible NOS1 expression and down-regulates ASCT1, a receptor for Syncytin-1, which is highly expressed in glia of individuals affected by multiple sclerosis. CREB3L1 is localized to the ER membrane until the ER undergoes stress, at which point CREB3L1 is cleaved sequentially by proteases SKI-1 and S2P and its N-terminus translocates into the nucleus. There are two isoforms of CREB3L1 that are produced as a result of alternative splicing events.

REFERENCES

- Omori, Y., et al. 2001. CREB-H: a novel mammalian transcription factor belonging to the CREB/ATF family and functioning via the box-B element with a liver-specific expression. *Nucleic Acids Res.* 29: 2154-2162.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611998. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Newman, J.R. and Keating, A.E. 2003. Comprehensive identification of human bZIP interactions with coiled-coil arrays. *Science* 300: 2097-2101.
- Chin, K.T., et al. 2005. The liver-enriched transcription factor CREB-H is a growth suppressor protein underexpressed in hepatocellular carcinoma. *Nucleic Acids Res.* 33: 1859-1873.
- Zhang, K., et al. 2006. Endoplasmic reticulum stress activates cleavage of CREBH to induce a systemic inflammatory response. *Cell* 124: 587-599.
- Panagopoulos, I., et al. 2007. Characterization of the native CREB3L2 transcription factor and the FUS/CREB3L2 chimera. *Genes Chromosomes Cancer* 46: 181-191.
- Bailey, D., et al. 2007. Trafficking of the bZIP transmembrane transcription factor CREB-H into alternate pathways of ERAD and stress-regulated intramembrane proteolysis. *Traffic* 8: 1796-1814.
- Schubert, S.W., et al. 2008. bZIP-Type transcription factors CREB and OASIS bind and stimulate the promoter of the mammalian transcription factor GCMa/Gcm1 in trophoblast cells. *Nucleic Acids Res.* 36: 3834-3846.

CHROMOSOMAL LOCATION

Genetic locus: CREB3L1 (human) mapping to 11p11.2.

SOURCE

CREB3L1 (F-8) is a mouse monoclonal antibody raised against amino acids 91-207 mapping near the N-terminus of CREB3L1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CREB3L1 (F-8) is recommended for detection of CREB3L1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CREB3L1 siRNA (h): sc-72995, CREB3L1 shRNA Plasmid (h): sc-72995-SH and CREB3L1 shRNA (h) Lentiviral Particles: sc-72995-V.


Molecular Weight of CREB3L1: 57 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CREB3L1 (F-8): sc-514635. Western blot analysis of CREB3L1 expression in Hep G2 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.