

# ASIC4 (E-8): sc-514818

## BACKGROUND

Degenerin/epithelial sodium channel (DEG/ENaC) superfamily members are amiloride-sensitive sodium channels that contain intracellular N- and C-termini, two hydrophobic transmembrane regions and a cysteine-containing extracellular loop. Acid sensing ion channel ASIC1, also designated ACCN2, BNAC2 and ASIC1 a, is present in brain as a 4.3-kb transcript with localization to rat dorsal root ganglia. *In situ* hybridization of rat brain suggests that ASIC1 is most abundant in the main olfactory bulb, cerebral cortex, hippocampal formation, habenula, basolateral amygdaloid nuclei and cerebellum. ASIC1 and H<sup>+</sup>-gated currents may contribute to the development of fear and anxiety. ASIC2, also designated ACCN1 (amiloride-sensitive cation channel 1, neuronal), mammalian degenerin, BNAC1 (MDEG) and brain Na<sup>+</sup> channel 1 mediate the normal detection of light touch. ASIC2 mRNA is abundant in brain, specifically in neurons. ASIC2 is expressed as 2.7- and 3.7-kb transcripts in brain and spinal cord tissues. ASIC3, also designated SLNAC1 and TNaC1, mediates detection of lasting pH changes and is involved in modulating moderate- to high-intensity pain sensation. ASIC4, also designated ACCN4 and BNAC4, is abundant in pituitary gland and is also present in the inner ear.

## REFERENCES

1. Garcia-Anoveros, J., et al. 1997. BNAC1 and BNAC2 constitute a new family of human neuronal sodium channels related to degenerins and epithelial sodium channels. *Proc. Natl. Acad. Sci. USA* 94: 1459-1464.
2. Waldmann, R., et al. 1997. A proton-gated cation channel involved in acid-sensing. *Nature* 386: 173-177.
3. Price, M.P., et al. 2000. The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature* 407: 1007-1011.

## CHROMOSOMAL LOCATION

Genetic locus: ASIC4 (human) mapping to 2q35; Asic4 (mouse) mapping to 1 C4.

## SOURCE

ASIC4 (E-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 1-24 at the N-terminus of ASIC4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ASIC4 (E-8) is available conjugated to agarose (sc-514818 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514818 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514818 PE), fluorescein (sc-514818 FITC), Alexa Fluor® 488 (sc-514818 AF488), Alexa Fluor® 546 (sc-514818 AF546), Alexa Fluor® 594 (sc-514818 AF594) or Alexa Fluor® 647 (sc-514818 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-514818 AF680) or Alexa Fluor® 790 (sc-514818 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-514818 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

ASIC4 (E-8) is recommended for detection of ASIC4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ASIC4 siRNA (h): sc-42411, ASIC4 siRNA (m): sc-42412, ASIC4 shRNA Plasmid (h): sc-42411-SH, ASIC4 shRNA Plasmid (m): sc-42412-SH, ASIC4 shRNA (h) Lentiviral Particles: sc-42411-V and ASIC4 shRNA (m) Lentiviral Particles: sc-42412-V.


Molecular Weight of ASIC4: 67 kDa.

Positive Controls: ASIC4 (h4): 293T Lysate: sc-111050, Hep G2 cell lysate: sc-2227 or K-562 whole cell lysate: sc-2203.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



ASIC4 (E-8): sc-514818. Western blot analysis of ASIC4 expression in K-562 (A), A-375 (B), HEL 92.1.7 (C), 3T3-L1 (D), RPE-J (E) and C6 (F) whole cell lysates.



ASIC4 (E-8): sc-514818. Western blot analysis of ASIC4 expression in non-transfected 293T: sc-117752 (A), human ASIC4 transfected 293T: sc-111050 (B), Hep G2 (C) and K-562 (D) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA