

IL-17R (C-12): sc-515358

BACKGROUND

Cytokines are small, soluble proteins with pleiotropic effects on a variety of cell types. Cytokines have a regulatory function over the immune system and mediate aspects of inflammatory response. They exert their biological effects through the binding of membrane-bound receptors which, in turn, initiate signal transduction cascades and elicit physiological changes in their target cell. Interleukin-17 (IL-17) and its cognate receptor, IL-17R, are an example of such a cytokine receptor pair. Originally identified as a rodent cDNA termed CTLA8, IL-17 is capable of inducing the secretion of IL-6 and IL-8 and augmenting the expression of ICAM-1 in human fibroblast cultures. The IL-17 protein exhibits a striking degree of homology with the HSV13 protein which mimics its function. The IL-17 receptor is a type I transmembrane protein 864 amino acids in length, that is highly expressed in spleen and kidney.

REFERENCES

1. Rouvier, E., et al. 1993. CTLA-8, cloned from an activated T cell, bearing AU-rich messenger RNA instability sequences, and homologous to a herpesvirus saimiri gene. *J. Immunol.* 150: 5445-5456.
2. Arend, W.P., et al. 1994. Binding of IL-1 α , IL-1 β , and IL-1 receptor antagonist by soluble IL-1 receptors and levels of soluble IL-1 receptors in synovial fluids. *J. Immunol.* 153: 4766-4774.
3. Yao, Z., et al. 1995. Herpesvirus saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 3: 811-821.
4. Okamura, H., et al. 1995. Cloning of a new cytokine that induces IFN- γ production by T cells. *Nature* 378: 88-91.
5. Yao, Z., et al. 1995. Human IL-17: a novel cytokine derived from T cells. *J. Immunol.* 155: 5483-5486.
6. Ihle, J.N. 1996. Janus kinases in cytokine signalling. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 351: 159-166.
7. Yao, Z., et al. 1996. Complete nucleotide sequence of the mouse CTLA8 gene. *Gene* 168: 223-225.

CHROMOSOMAL LOCATION

Genetic locus: Il17ra (mouse) mapping to 6 F1.

SOURCE

IL-17R (C-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 841-864 at the C-terminus of IL-17R of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-515358 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

IL-17R (C-12) is recommended for detection of IL-17R of mouse origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-17R siRNA (m): sc-40038, IL-17R shRNA Plasmid (m): sc-40038-SH and IL-17R shRNA (m) Lentiviral Particles: sc-40038-V.

Molecular Weight of IL-17R: 120 kDa.


Positive Controls: IL-17R (m2): 293T Lysate: sc-178795.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



IL-17R (C-12): sc-515358. Western blot analysis of IL-17R expression in non-transfected: sc-117752 (A) and mouse IL-17R transfected: sc-178795 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.