

BMP-8 (H-4): sc-515399

BACKGROUND

Bone morphogenic proteins (BMPs) are members of the TGF β superfamily. BMPs are involved in the induction of cartilage and bone formation. *In vivo* studies have shown that BMP-2 (also designated BMP-2A) and BMP-3 can independently induce cartilage formation. Smad3 association with the TGF β receptor complex and Smad1 translocation to the nucleus are observed after the addition of BMP-4 (also designated BMP-2B), suggesting that BMP-4 may play a role in activation of the Smad pathway. BMP-5, BMP-6 and BMP-7 all share high sequence homology with BMP-2, indicating that they each may be able to induce cartilage formation. BMP-8 is thought to be involved in early development, as detectable expression has not been found in adult organs. Two BMP-8 proteins exist, namely BMP-8A and BMP-8B (also designated OP-2), and are encoded by two distinct genes.

REFERENCES

1. Wozney, J.M., et al. 1988. Novel regulators of bone formation: molecular clones and activities. *Science* 242: 1528-1534.
2. Massague, J. 1990. The transforming growth factor- β family. *Annu. Rev. Cell Biol.* 6: 597-641.
3. Celeste, A.J., et al. 1990. Identification of transforming growth factor β family members present in bone-inductive protein purified from bovine bone. *Proc. Natl. Acad. Sci. USA* 87: 9843-9847.
4. Oskaynak, E., et al. 1992. Osteogenic protein-2. A new member of the transforming growth factor- β superfamily expressed early in embryogenesis. *J. Biol. Chem.* 267: 25220-25227.

CHROMOSOMAL LOCATION

Genetic locus: BMP8A (human) mapping to 1p34.3, BMP8B (human) mapping to 1p34.2; Bmp8b (mouse) mapping to 4 D2.2.

SOURCE

BMP-8 (H-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 265-287 at the N-terminus of BMP-8B of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

BMP-8 (H-4) is available conjugated to agarose (sc-515399 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515399 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515399 PE), fluorescein (sc-515399 FITC), Alexa Fluor[®] 488 (sc-515399 AF488), Alexa Fluor[®] 546 (sc-515399 AF546), Alexa Fluor[®] 594 (sc-515399 AF594) or Alexa Fluor[®] 647 (sc-515399 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-515399 AF680) or Alexa Fluor[®] 790 (sc-515399 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-515399 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

BMP-8 (H-4) is recommended for detection of precursor and mature BMP-8A and BMP-8B of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of BMP-8: 45 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Raji whole cell lysate: sc-364236 or NCI-H1299 whole cell lysate: sc-364234.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



BMP-8 (H-4): sc-515399. Western blot analysis of BMP-8 expression in HeLa (A), NCI-H1299 (B) and Raji (C) whole cell lysates.



BMP-8 (H-4): sc-515399. Western blot analysis of BMP-8 expression in c4 (A), Neuro-2A (B) and BW5147 (C) whole cell lysates.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.