


IL-1 β (H-8): sc-515786



The Power to Question

BACKGROUND

Two forms of interleukin-1, designated IL-1 α and IL-1 β , have been described. Although encoded by distinct genes and exhibiting roughly only 25% sequence identity, IL-1 α and IL-1 β bind to the same receptor and seem to elicit similar biological responses. IL-1 production is generally thought to be associated with inflammation, but it has also been shown to be expressed during kidney development, thymocyte differentiation and cartilage degradation. IL-1 plays a critical role in the regulation of immune response and inflammation, acting as an activator of T and B lymphocytes and natural killer (NK) cells. In T cells, IL-1 stimulates the production of IL-2 and selectively inhibits IL-4 expression. IL-1 induces B cell proliferation and maturation, and immunoglobulin synthesis. NK cells require IL-1 β for production of the anti-pathogen IFN- γ . IL-1 has also been implicated in several pathological conditions including rheumatoid arthritis, inflammatory bowel disease and atherosclerosis.

REFERENCES

1. Auron, P.E., et al. 1985. Nucleotide sequence of human monocyte interleukin-1 precursor cDNA. Proc. Natl. Acad. Sci. USA 81: 7907-7911.
2. March, C.J., et al. 1985. Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs. Nature 315: 641-647.
3. Dinarello, C.A. 1991. Interleukin-1 and interleukin-1 antagonism. Blood 77: 1627-1652.
4. Sadouk, M.B., et al. 1995. Human synovial fibroblasts coexpress IL-1 receptor type I and type II mRNA. The increased level of the IL-1 receptor in osteoarthritis cells is related to an increased level of the type I receptor. Lab. Invest. 73: 347-355.
5. Lonnemann, G., et al. 1995. Cytokines in human renal interstitial fibrosis. I. Interleukin-1 is a paracrine growth factor for cultured fibrosis-derived kidney fibroblasts. Kidney Int. 47: 837-844.
6. Zúñiga-Pflücker, J.C., et al. 1995. Requirement for TNF α and IL-1 α in fetal thymocyte commitment and differentiation. Science 268: 1906-1909.
7. Sandborg, C.I., et al. 1995. IL-4 expression in human T cells is selectively inhibited by IL-1 α and IL-1 β . J. Immunol. 155: 5206-5212.
8. Leung, L.Y., et al. 2007. A new microrheometric approach reveals individual and cooperative roles for TGF β 1 and IL-1 β in fibroblast-mediated stiffening of collagen gels. FASEB J. 21: 2064-2073.

CHROMOSOMAL LOCATION

Genetic locus: IL1B (human) mapping to 2q14.1.

SOURCE

IL-1 β (H-8) is a mouse monoclonal antibody raised against amino acids 117-269 of IL-1 β of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

IL-1 β (H-8) is recommended for detection of IL-1 β of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-1 β siRNA (h): sc-39615, IL-1 β shRNA Plasmid (h): sc-39615-SH and IL-1 β shRNA (h) Lentiviral Particles: sc-39615-V.

Molecular Weight of IL-1 β precursor: 31 kDa.


Molecular Weight of mature IL-1 β : 17 kDa.

Positive Controls: IL-1 β (h): 293 Lysate: sc-111184 or HT-1080 whole cell lysate: sc-364183.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



IL-1 β (H-8): sc-515786. Western blot analysis of IL-1 β expression in non-transfected: sc-110760 (A) and human IL-1 β transfected: sc-111184 (B) 293 whole cell lysates.



IL-1 β (H-8): sc-515786. Western blot analysis of IL-1 β expression in HT-1080 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.