

mouse anti-goat IgG-CFL 405: sc-516240

BACKGROUND

Santa Cruz Biotechnology offers a wide variety of control immunoglobulin and control sera for a large selection of species, including mouse, rabbit, goat, chicken, rat, hamster, canine, guinea pig and sheep. Control immunoglobulin and immunoglobulin conjugates are useful negative controls. Normal sera is offered to be used as blocking reagents. Santa Cruz Biotechnology offers affinity purified normal immunoglobulins and immunoglobulin conjugates for use as negative controls in applications including flow cytometry, immunohistochemistry, immunofluorescence, Western Blotting and immunoprecipitation. Agarose (AC) conjugated IgGs are provided for immunoprecipitation; horseradish peroxidase (HRP) conjugates are provided for Western Blotting and immunohistochemistry; and Biotin (B) conjugates are provided for immunohistochemistry. A broad range of fluorescent conjugated controls are also available for use in flow cytometry and immunofluorescence applications. Most control immunoglobulins are available as unconjugated controls or as FITC (fluorescein isothiocyanate), PE (phycoerythrin), PE-Cy5 (phycoerythrin-Cy5), PE-Cy7 (phycoerythrin-Cy7), APC (allophycocyanin) and APC-Cy7 (allophycocyanin-Cy7) conjugates. Additional conjugates include Alexa Fluor® 488, Alexa Fluor® 647, Alexa Fluor® 405, PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex-Cy 5.5). Isotype specific control immunoglobulins include classes such as mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, IgM and IgA, rat IgG₁, IgG_{2a}, IgG_{2b} and IgM, Armenian hamster IgG, and both goat and rabbit IgG.

SOURCE

mouse anti-goat IgG-CFL 405 is a pre-adsorbed, affinity purified secondary antibody raised in mouse against goat IgG and conjugated to CruzFluor™ 405 (CFL 405).

PRODUCT

Each vial contains 200 µg mouse IgG (pre-adsorbed with human IgG) in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

APPLICATIONS

mouse anti-goat IgG-CFL 405 is recommended for detection of goat IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400), immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400) and flow cytometry (0.5-1 µg per 1 x 10⁶ cells) of immunoglobulin-rich tissues and cells. Optimal dilution to be determined by titration.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Texas Red® and Alexa Fluor® are registered trademarks of Molecular Probes, Inc., Oregon, USA

RECOMMENDED SUPPORT PRODUCTS

A. TISSUE CULTURE CELLS

- CrystalCruz™ Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336

B. FROZEN TISSUE SECTIONS

- Organo/Limonene Mount, non-toxic alternative to Permount, 100 ml: sc-45087
- UltraCruz® Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with *in situ* hybridization, 30 ml: sc-45088

C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS

- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973