# SANTA CRUZ BIOTECHNOLOGY, INC.

# Thyroglobulin (2H11): sc-51708



### BACKGROUND

Thyroglobulin is a large preprotein containing multiple glycosylation sites. Located in the thyroid gland, thyroglobulin is the precursor of the iodinated thyroid hormones thyroxine and triiodothyronine. Thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum, prior to forming dimers. This dimerization, as well as export of Thyroglobulin to the Golgi complex, has been shown to require Ca<sup>2+</sup>. Defects in Thyroglobulin are known to cause some types of goiter (an enlargement of the thyroid gland). This condition is thought to result from defective dimerization and transport of thyroglobulin to the Golgi complex.

#### **REFERENCES**

- 1. Malthiery, Y. and Lissitzky, S. 1987. Primary structure of human Thyroglobulin deduced from the sequence of its 8448-base complementary DNA. Eur. J. Biochem. 165: 491-498.
- 2. Mallet, B., et al. 1995. N-glycans modulate in vivo and in vitro thyroid hormone synthesis. Study at the N-terminal domain of Thyroglobulin. J. Biol. Chem. 270: 29881-29888.
- 3. Prabakaran, D., et al. 1996. Oligomeric assembly of thrombospondin in the endoplasmic reticulum of thyroid epithelial cells. Eur. J. Cell Biol. 70: 134-141.

## **CHROMOSOMAL LOCATION**

Genetic locus: TG (human) mapping to 8g24.22; Tg (mouse) mapping to 15 D2.

### SOURCE

Thyroglobulin (2H11) is a mouse monoclonal antibody raised against human thyroid follicular cells.

## PRODUCT

Each vial contains 200  $\mu g$  IgG\_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

#### **APPLICATIONS**

Thyroglobulin (2H11) is recommended for detection of Thyroglobulin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffinembedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Thyroglobulin siRNA (h): sc-63346, Thyroglobulin siRNA (m): sc-63347, Thyroglobulin shRNA Plasmid (h): sc-63346-SH, Thyroglobulin shRNA Plasmid (m): sc-63347-SH, Thyroglobulin shRNA (h) Lentiviral Particles: sc-63346-V and Thyroglobulin shRNA (m) Lentiviral Particles: sc-63347-V.

Molecular Weight of Thyroglobulin isoforms: 305/298 kDa.

Positive Controls: human thyroid extract: sc-363782.

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-lgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

#### DATA



Thyroglobulin (2H11): sc-51708. Western blot analysis of Thyroglobulin expression in human thyroid tissue extract

#### **SELECT PRODUCT CITATIONS**

- 1. Linda Messina, R., et al. 2012. Reactivation of P53 mutants by prima-1 in thyroid cancer cells. Int. J. Cancer 130: 2259-2270.
- 2. Gianì, F., et al. 2015. Thyrospheres from normal or malignant thyroid tissue have different biological, functional, and genetic features. J. Clin. Endocrinol. Metab. 100: E1168-E1178.
- 3. Tang, D., et al. 2015. Apoptosis and anergy of T cell induced by pancreatic stellate cells-derived galectin-1 in pancreatic cancer. Tumour Biol. 36: 5617-5626.
- 4. Sun, Z., et al. 2019. PTEN-knockdown disrupts the morphology, growth pattern and function of Nthy-Ori 3-1 cells by downregulating PAX8 expression. Oncol. Lett. 18: 6732-6740.
- 5. Hassan, N.H. and Amin, M.A. 2022. Resveratrol thyro-protective role in fluorosis rat model (histo-morphometric, biochemical and ultrastructural study). Tissue Cell 80: 101986.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.