

CHRDL1 (1D10): sc-517585

BACKGROUND

The patterning of the central nervous system (CNS) relies on the interaction of multiple signaling molecules, including Shh (sonic hedgehog homolog), Wnt proteins and BMPs (bone morphogenetic proteins), with their antagonists, namely chordin and Noggin. Chordin is a key developmental protein that dorsalizes early vertebrate embryonic tissues by binding to ventralizing TGF- β -like BMPs and sequestering them in latent complexes. CHRDL1 (Chordin-like protein 1), also known as CHL, VOPT (Ventroptin) or NRLN1 (Neuralin-1), is a 450 amino acid secreted protein that contains three VWFC domains. Functioning in a similar manner to chordin, CHRDL1 binds to BMP-4 and prevents the interaction of BMP-4 with its target receptors, thereby antagonizing BMP-4 activity. Additionally, CHRDL1 is thought to play a role in dorsoventral axis formation, embryonic bone formation and angiogenesis, as well as in the differentiation of neural stem cells.

REFERENCES

1. Coffinier, C., Tran, U., Larraín, J. and De Robertis, E.M. 2001. Neuralin-1 is a novel Chordin-related molecule expressed in the mouse neural plate. *Mech. Dev.* 100: 119-122.
2. Sakuta, H., Suzuki, R., Takahashi, H., Kato, A., Shintani, T., Iemura, S., Ueno, N. and Noda, M. 2001. Ventroptin: a BMP-4 antagonist expressed in a double-gradient pattern in the retina. *Science* 293: 111-115.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300350. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Chimal-Monroy, J., Rodriguez-Leon, J., Montero, J.A., Gañan, Y., Macias, D., Merino, R. and Hurler, J.M. 2003. Analysis of the molecular cascade responsible for mesodermal limb chondrogenesis: Sox genes and BMP signaling. *Dev. Biol.* 257: 292-301.
5. Chen, D., Zhao, M. and Mundy, G.R. 2004. Bone morphogenetic proteins. *Growth Factors* 22: 233-241.
6. Gazzero, E. and Canalis, E. 2006. Bone morphogenetic proteins and their antagonists. *Rev. Endocr. Metab. Disord.* 7: 51-65.
7. Kosinski, C., Li, V.S., Chan, A.S., Zhang, J., Ho, C., Tsui, W.Y., Chan, T.L., Mifflin, R.C., Powell, D.W., Yuen, S.T., Leung, S.Y. and Chen, X. 2007. Gene expression patterns of human colon tops and basal crypts and BMP antagonists as intestinal stem cell niche factors. *Proc. Natl. Acad. Sci. USA* 104: 15418-15423.
8. Kane, R., Godson, C. and O'Brien, C. 2008. Chordin-like 1, a bone morphogenetic protein-4 antagonist, is upregulated by hypoxia in human retinal pericytes and plays a role in regulating angiogenesis. *Mol. Vis.* 14: 1138-1148.

CHROMOSOMAL LOCATION

Genetic locus: CHRDL1 (human) mapping to Xq23.

SOURCE

CHRDL1 (1D10) is a mouse monoclonal antibody raised against recombinant GST-tagged CHRDL1 of human origin.

PRODUCT

Each vial contains 50 μ g IgG₁ kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CHRDL1 (1D10) is recommended for detection of CHRDL1 of human origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CHRDL1 siRNA (h): sc-91247, CHRDL1 shRNA Plasmid (h): sc-91247-SH and CHRDL1 shRNA (h) Lentiviral Particles: sc-91247-V.

Molecular Weight of CHRDL1: 60 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.