

β5 Tubulin (87CT59.3.7): sc-517663

BACKGROUND

Tubulin is a major cytoskeleton component that has five distinct forms, designated α , β , γ , δ and ϵ Tubulin. α and β Tubulins form heterodimers which multimerize to form a microtubule filament. Multiple β Tubulin isoforms ($\beta 1$, $\beta 2$, $\beta 3$, $\beta 4$, $\beta 5$, $\beta 6$ and $\beta 8$) have been characterized and are expressed in mammalian tissues. $\beta 1$ and $\beta 4$ are present throughout the cytosol, $\beta 2$ is present in the nuclei and nucleoplasm, and $\beta 3$ is a neuron-specific cytoskeletal protein. γ Tubulin forms the gammaosome, which is required for nucleating microtubule filaments at the centrosome. Both δ Tubulin and ϵ Tubulin are associated with the centrosome. δ Tubulin is a homolog of the *Chlamydomonas* δ Tubulin Uni3 and is found in association with the centrioles, whereas ϵ Tubulin localizes to the pericentriolar material. ϵ Tubulin exhibits a cell-cycle-specific pattern of localization, first associating with only the older of the centrosomes in a newly duplicated pair and later associating with both centrosomes.

REFERENCES

- Weisenberg, R. 1981. Invited review: the role of nucleotide triphosphate in Actin and Tubulin assembly and function. *Cell Motil.* 1: 485-497.
- Burns, R.G. 1991. α -, β -, and γ -Tubulins: sequence comparisons and structural constraints. *Cell Motil. Cytoskeleton* 20: 181-189.
- Zheng, Y., et al. 1991. γ Tubulin is present in *Drosophila melanogaster* and *Homo sapiens* and is associated with the centrosome. *Cell* 65: 817-823.
- Leask, A. and Stearns, T. 1998. Expression of amino- and carboxyl-terminal γ and β Tubulin mutants in cultured epithelial cells. *J. Biol. Chem.* 273: 2661-2668.
- Ludueno, R.F. 1998. Multiple forms of Tubulin: different gene products and covalent modifications. *Int. Rev. Cytol.* 178: 207-275.
- Walss, C., et al. 1999. Presence of the $\beta 2$ isotype of Tubulin in the nuclei of cultured mesangial cells from rat kidney. *Cell Motil. Cytoskeleton* 42: 274-284.
- Modig, C., et al. 1999. Identification of $\beta 3$ and $\beta 4$ Tubulin isotypes in cold-adapted microtubules from Atlantic cod (*Gadus morhua*): antibody mapping and cDNA sequencing. *Cell Motil. Cytoskeleton* 42: 315-330.
- Woulfe, J. and Munoz, D. 2000. Tubulin immunoreactive neuronal intranuclear inclusions in the human brain. *Neuropathol. Appl. Neurobiol.* 26: 161-171.
- Chang, P. and Stearns, T. 2000. δ Tubulin and ϵ Tubulin: two new human centrosomal tubulins reveal new aspects of centrosome structure and function. *Nat. Cell Biol.* 2: 30-35.

CHROMOSOMAL LOCATION

Genetic locus: TUBB (human) mapping to 6p21.33; Tubb5 (mouse) mapping to 17 B1.

SOURCE

$\beta 5$ Tubulin (87CT59.3.7) is a mouse monoclonal antibody raised against recombinant $\beta 5$ Tubulin of human origin.

PRODUCT

Each vial contains 100 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

$\beta 5$ Tubulin (87CT59.3.7) is recommended for detection of $\beta 5$ Tubulin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for $\beta 5$ Tubulin siRNA (h): sc-106649, $\beta 5$ Tubulin siRNA (m): sc-140335, $\beta 5$ Tubulin shRNA Plasmid (h): sc-106649-SH, $\beta 5$ Tubulin shRNA Plasmid (m): sc-140335-SH, $\beta 5$ Tubulin shRNA (h) Lentiviral Particles: sc-106649-V and $\beta 5$ Tubulin shRNA (m) Lentiviral Particles: sc-140335-V.

Molecular Weight of $\beta 5$ Tubulin: 55 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 3) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.