

# NHE-1 (C-7): sc-518042

## BACKGROUND

Na<sup>+</sup>/H<sup>+</sup> exchangers 1–6 (Na<sup>+</sup>/H<sup>+</sup> antiporters, NHE-1-6) are integral membrane proteins that are expressed in most mammalian tissues, where they regulate intracellular pH and cell volume. NHEs mediate the secondary active extrusion of hydrogen (H<sup>+</sup>) ions out of cells in exchange for extracellular sodium (Na<sup>+</sup>). Excluding NHE-1, which is ubiquitously expressed, the NHE isoforms NHE-2–6 have distinct tissue- and cell type-dependent expression and inhibitory characteristics by amiloride analogs. Human NHE-1 protein, known also as solute carrier family 9 isoform-1 (SLC9A1), is a ten transmembrane domain-spanning receptor that contains an N-terminal amphiphatic domain and two putative N-glycosylation sites.

## REFERENCES

- Sardet, C., et al. 1989. Molecular cloning, primary structure, and expression of the human growth factor-activatable Na<sup>+</sup>/H<sup>+</sup> antiporter. *Cell* 56: 271-280.
- Orlowski, J., et al. 1992. Molecular cloning of putative members of the Na/H exchanger gene family. cDNA cloning, deduced amino acid sequence, and mRNA tissue expression of the rat Na/H exchanger NHE-1 and two structurally related proteins. *J. Biol. Chem.* 267: 9331-9339.
- Fliegel, L., et al. 1993. Cloning and analysis of the human myocardial Na<sup>+</sup>/H<sup>+</sup> exchanger. *Mol. Cell. Biochem.* 125: 137-143.
- Biemesderfer, D., et al. 1993. NHE-3: a Na<sup>+</sup>/H<sup>+</sup> exchanger isoform of renal brush border. *Am. J. Physiol.* 265: 736-742.
- Noel, J. and Pouyssegur, J. 1995. Hormonal regulation, pharmacology, and membrane sorting of vertebrate Na<sup>+</sup>/H<sup>+</sup> exchanger isoforms. *Am. J. Physiol.* 268: 283-296.
- Klanke, C.A., et al. 1995. Molecular cloning and physical and genetic mapping of a novel human Na<sup>+</sup>/H<sup>+</sup> exchanger (NHE-5/SLC9A5) to chromosome 16q22.1. *Genomics* 25: 615-622.
- Cox, G.A., et al. 1997. Sodium/hydrogen exchanger gene defect in slow-wave epilepsy mutant mice. *Cell* 91: 139-148.

## CHROMOSOMAL LOCATION

Genetic locus: SLC9A1 (human) mapping to 1p36.11; Slc9a1 (mouse) mapping to 4 D2.3.

## SOURCE

NHE-1 (C-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 42-61 near the N-terminus of NHE-1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

NHE-1 (C-7) is recommended for detection of NHE-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NHE-1 siRNA (h): sc-42650, NHE-1 siRNA (m): sc-42651, NHE-1 shRNA Plasmid (h): sc-42650-SH, NHE-1 shRNA Plasmid (m): sc-42651-SH, NHE-1 shRNA (h) Lentiviral Particles: sc-42650-V and NHE-1 shRNA (m) Lentiviral Particles: sc-42651-V.

Molecular Weight of NHE-1 precursor: 90 kDa.

Molecular Weight of glycosylated NHE-1: 110-130 kDa.


Molecular Weight of NHE-1 dimer: 210 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, HeLa whole cell lysate: sc-2200 or MOLT-4 cell lysate: sc-2233.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



NHE-1 (C-7): sc-518042. Western blot analysis of NHE-1 expression in non-transfected (A) and mouse NHE-1 transfected (B), K-562 (C), HeLa (D), MOLT-4 (E) and Caki-1 (F) whole cell lysates.



NHE-1 (C-7): sc-518042. Western blot analysis of NHE-1 expression in MCF7 whole cell lysate. Detection reagent used: m-IgG Fc BP-HRP: sc-525409.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.