SANTA CRUZ BIOTECHNOLOGY, INC.

NGF (B-4): sc-518166



BACKGROUND

Neurotrophins function to regulate naturally occurring cell death of neurons during development. The prototype neurotrophin is nerve growth factor (NGF), originally discovered in the 1950s as a soluble peptide promoting the survival of, and neurite outgrowth from, sympathetic ganglia. Three additional structurally homologous neurotrophic factors have been identified. These include brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4) (also designated NT-5). These various neurotrophins stimulate the *in vitro* survival of distinct, but partially overlapping, populations of neurons. The cell surface receptors through which neurotrophins mediate their activity have been identified. For instance, the Trk A receptor is the preferential receptor for NGF, but also binds NT-3 and NT-4. The Trk B receptor binds both BDNF and NT-4 equally well, and binds NT-3 to a lesser extent, while the Trk C receptor only binds NT-3.

REFERENCES

- Oppenhiem, R.W. 1991. Cell death during development of the nervous system. Annu. Rev. Neurosci. 14: 453-501.
- 2. Thoenen, H. 1991. The changing scene of neurotrophic factors. Trends Neurosci. 14: 165-170.
- Chao, M.V. 1992. Neurotrophin receptors: a window into neuronal differentiation. Neuron 9: 583-593.
- Korsching, S. 1993. The neurotrophic factor concept: a reexamination. J. Neurosci. 13: 2739-2748.
- 5. lp, N.Y., et al. 1993. Similarities and differences in the way neurotrophins interact with the Trk receptors in neuronal and nonneuronal cells. Neuron 10: 137-149.
- 6. Klein, R. 1994. Role of neurotrophins in mouse neuronal development. FASEB J. 8: 738-744.

CHROMOSOMAL LOCATION

Genetic locus: NGF (human) mapping to 1p13.2.

SOURCE

NGF (B-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 118-146 at the N-terminus of NGF of human origin.

PRODUCT

Each vial contains 200 μg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NGF (B-4) is available conjugated to agarose (sc-518166 AC), 500 µg/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-518166 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518166 PE), fluorescein (sc-518166 FITC), Alexa Fluor[®] 488 (sc-518166 AF488), Alexa Fluor[®] 546 (sc-518166 AF546), Alexa Fluor[®] 594 (sc-518166 AF594) or Alexa Fluor[®] 647 (sc-518166 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-518166 AF680) or Alexa Fluor[®] 790 (sc-518166 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

NGF (B-4) is recommended for detection of NGF precursor and mature forms of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NGF siRNA (h): sc-43970, NGF shRNA Plasmid (h): sc-43970-SH and NGF shRNA (h) Lentiviral Particles: sc-43970-V.

Molecular Weight of mature NGF: 13 kDa.

Molecular Weight of NGF precursor: 27 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



human recombinant NGF. Detection reagent used: mlqG κ BP-HRP: sc-516102.

SELECT PRODUCT CITATIONS

 Abbastabar, M., et al. 2023. Expression status of Rap1 pathway-related genes in liver metastases compared with corresponding primary colorectal cancer. Cancers 16: 171.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.