



## CS protein (PVC-2): sc-51855

### BACKGROUND

The circumsporozoite (CS) protein is a cell surface protein of the sporozoite, the stage of the life cycle of malaria parasites (*Plasmodium* spp.) that infects the vertebrate host. The sporozoite stage of the *Plasmodium* parasite is formed by budding from a multinucleate oocyst in the mosquito midgut. Sporozoites must infect the salivary glands of the mosquito vector and the liver of the mammalian host; both events depend on the CS protein. The amounts of CS protein correlate with the extent of development of the inner membranes and associated microtubules underneath the oocyst outer membrane, which normally demarcate focal budding sites.

### REFERENCES

- Romero, P., Maryanski, J.L., Corradin, G., Nussenzweig, R.S., Nussenzweig, V. and Zavala, F. 1989. Cloned cytotoxic T cells recognize an epitope in the circumsporozoite protein and protect against malaria. *Nature* 341: 323-326.
- Greenwood, J., Willis, A.E. and Perham, R.N. 1991. Multiple display of foreign peptides on a filamentous bacteriophage. Peptides from *Plasmodium falciparum* circumsporozoite protein as antigens. *J. Mol. Biol.* 220: 821-827.
- Hughes, A.L. 1991. Circumsporozoite protein genes of malaria parasites (*Plasmodium* spp.): evidence for positive selection on immunogenic regions. *Genetics* 127: 345-353.
- Cerami, C., Frevert, U., Sinnis, P., Takacs, B., Clavijo, P., Santos, M.J. and Nussenzweig, V. 1992. The basolateral domain of the hepatocyte plasma membrane bears receptors for the circumsporozoite protein of *Plasmodium falciparum* sporozoites. *Cell* 70: 1021-1033.
- Sedegah, M., Hedstrom, R., Hobart, P. and Hoffman, S.L. 1994. Protection against malaria by immunization with plasmid DNA encoding circumsporozoite protein. *Proc. Natl. Acad. Sci. USA* 91: 9866-9870.
- Menard, R., Sultan, A.A., Cortes, C., Altszuler, R., van Dijk, M.R., Janse, C.J., Waters, A.P., Nussenzweig, R.S. and Nussenzweig, V. 1997. Circumsporozoite protein is required for development of malaria sporozoites in mosquitoes. *Nature* 385: 336-340.
- Thathy, V., Fujioka, H., Gantt, S., Nussenzweig, R., Nussenzweig, V. and Menard, R. 2002. Levels of circumsporozoite protein in the *Plasmodium* oocyst determine sporozoite morphology. *EMBO J.* 21: 1586-1596.

### SOURCE

CS protein (PVC-2) is a mouse monoclonal antibody raised against recombinant CS protein of *Plasmodium vivax* origin.

### PRODUCT

Each vial contains 100 µg IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### APPLICATIONS

CS protein (PVC-2) is recommended for detection of CS protein of *Plasmodium vivax* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

Molecular Weight of CS: 54 kDa.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.