

# IL-2 (LNKB-2): sc-52017

## BACKGROUND

Lymphokines are a group of signaling molecules involved in communication between cells of the immune system. Lymphokines secreted by activated lymphocytes include proteins such as interleukin-2 (IL-2). This protein is secreted primarily by helper T cells that have been activated through the T cell receptor complex or by other mitogens. Cells targeted by IL-2 include activated T helper and cytotoxic T cells, inducing their proliferation. The secretion of IL-2 can also act as a growth factor for B cells. To date, three different IL-2-dependent signal transduction pathways have been identified: the c-Fos/c-Jun induction pathway mediated by Src family protein-tyrosine kinases, the c-Myc induction pathway and the Rapamycin-sensitive pathway, all of which result in the induction of Bcl-2. In addition, the transcription factor NFAT has been shown to play a major role in the regulation of IL-2 transcription and correlates to an age-related decline in the expression of IL-2.

## REFERENCES

- Smith, K.A. 1980. T cell growth factor. *Immunol. Rev.* 51: 337-357.
- Taniguchi, T., et al. 1983. Structure and expression of a cloned cDNA for human interleukin-2. *Nature* 302: 305-310.
- Lowenthal, J.W., et al. 1985. Similarities between interleukin-2 receptor number and affinity on activated B and T lymphocytes. *Nature* 315: 669-672.
- Guy, G.R., et al. 1990. Lymphokine signal transduction. *Prog. Growth Factor Res.* 2: 45-70.
- Germann, T., et al. 1991. Components of an antigen-/T cell receptor-independent pathway of lymphokine production. *Eur. J. Immunol.* 21: 1857-1861.
- Miyazaki, T., et al. 1995. Three distinct IL-2 signaling pathways mediated by Bcl-2, c-Myc and Lck cooperate in hematopoietic cell proliferation. *Cell* 81: 223-231.
- Eljaafari, A., et al. 1995. Contribution of p56Lck to the upregulation of cytokine production and T cell proliferation by IL-2 in human CD3-stimulated T cell clones. *Cell. Immunol.* 160: 152-156.
- Pahlavani, M.A., et al. 1995. The age-related decline in the induction of IL-2 transcription is correlated to changes in the transcription factor NFAT. *Cell. Immunol.* 165: 84-91.

## CHROMOSOMAL LOCATION

Genetic locus: IL2 (human) mapping to 4q27.

## SOURCE

IL-2 (LNKB-2) is a mouse monoclonal antibody raised against recombinant IL-2 of human origin.

## PRODUCT

Each vial contains 100 µg IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS


IL-2 (LNKB-2) is recommended for detection of IL-2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for IL-2 siRNA (h): sc-39619, IL-2 shRNA Plasmid (h): sc-39619-SH and IL-2 shRNA (h) Lentiviral Particles: sc-39619-V.


Molecular Weight of IL-2: 15 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HuT 78 whole cell lysate: sc-2208 or CCRF-HSB-2 cell lysate: sc-2265.

## DATA



IL-2 (LNKB-2): sc-52017. Western blot analysis of IL-2 expression in Jurkat (A), CCRF-CEM (B), HuT 78 (C), H9 (D), SUP-T1 (E) and CCRF-HSB-2 (F) whole cell lysates.



IL-2 (LNKB-2): sc-52017. Western blot analysis of IL-2 expression in MOLT-4 whole cell lysate (A) and human spleen tissue extract (B).

## SELECT PRODUCT CITATIONS

- Burbelo, P.D., et al. 2015. Lack of evidence for molecular mimicry in HIV-infected subjects. *PLoS ONE* 10: e0127662.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.