



Listeria monocytogenes (LZH1): sc-52059

BACKGROUND

The genus *Listeria* includes the species *Listeria monocytogenes*, a gram-positive bacterium that is motile by means of flagella, though it can also move inside of eukaryotic cells by polymerizing actin comet tails with a protein called ActA. *L. monocytogenes* is the only species of the genus *Listeria* that has been clearly documented as a pathogen for humans. It is food-borne, intracellular, and interacts with infected host cells both prior to entry and during the intracellular phase of infection. Uptake by the host cell is stimulated by the binding of listerial internalins (InI) to host cell adhesion factors such as E-cadherin or Met, which then exploits the junction forming-apparatus of the host into internalizing the bacterium. Infection by *L. monocytogenes* causes the disease listeriosis, which can manifest into meningitis and septicemia. Listeriosis occurs mainly in newborn infants, elderly patients and patients who are immunocompromised.

REFERENCES

- Han, Y., Linton, R.H. and Nelson, P.E. 2004. Effects of recovery, plating, and inoculation methods on quantification of *Escherichia coli* O157:H7 and *Listeria monocytogenes* from strawberries. *J. Food Prot.* 67: 2436-2442.
- Gómez, N., García, D., Alvarez, I., Condón, S. and Raso, J. 2005. Modelling inactivation of *Listeria monocytogenes* by pulsed electric fields in media of different pH. *Int. J. Food Microbiol.* 103: 199-206.
- Munder, A., Zelmer, A., Schmiedl, A., Dittmar, K.E., Rohde, M., Dorsch, M., Otto, K., Hedrich, H.J., Tümmler, B., Weiss, S. and Tschernig, T. 2005. Murine pulmonary infection with *Listeria monocytogenes*: differential susceptibility of BALB/c, C57BL/6 and DBA/2 mice. *Microbes Infect.* 7: 600-611.
- Rodríguez-Lázaro, D., Pla, M., Scorti, M., Vázquez-Boland, J.A. and Monzó, H.J. 2005. A novel real-time PCR for *Listeria monocytogenes* that monitors analytical via an internal amplification control. *Appl. Environ. Microbiol.* 71: 9008-9012.
- Bozariis, I.S. and Nychas, G.J. 2006. Effect of nisin on growth boundaries of list various temperatures, pH and water activities. *Food Microbiol.* 23: 779-784.
- Kathariou, S., Graves, L., Buchrieser, C., Glaser, P., Siletsky, R.M. and Swaminathan, B. 2006. Involvement of closely related strains of a new clonal group of *Listeria monocytogenes* in the 1998-99 and 2002 multi-state outbreaks of foodborne listeriosis in the United States. *Foodborne Pathog. Dis.* 3: 292-302.
- Okada, Y., Okada, N., Makino, S., Asakura, H., Yamamoto, S. and Igimi, S. 2006. The sigma factor RpoN (σ 54) is involved in osmotolerance in *Listeria monocytogenes*. *FEMS Microbiol. Lett.* 263: 54-60.
- Ozören, N., Masumoto, J., Franchi, L., Kanneganti, T.D., Body-Malapel, M., Ertürk, I., Jagirdar, R., Zhu, L., Inohara, N., Bertin, J., Coyle, A., Grant, EP. and Núñez, G. 2006. Distinct roles of TLR2 and the adaptor ASC in IL-1 β /IL-18 to *Listeria monocytogenes*. *J. Immunol.* 176: 4337-4342.
- Rossmannith, P., Krassnig, M., Wagner, M. and Hein, I. 2006. Detection of *Listeria monocytogenes* in food using a combined enrichment/real-time PCR method targeting the prfA gene. *Res. Microbiol.* 157: 763-771.

SOURCE

Listeria monocytogenes (LZH1) is a mouse monoclonal antibody raised against *Listeria monocytogenes*.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Listeria monocytogenes (LZH1) is recommended for detection of intact cells as well as the outer membrane fraction of *Listeria monocytogenes* by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.