



Ricin B (RB127): sc-52191

BACKGROUND

Ricin, a type II ribosomal inactivating protein, inhibits protein biosynthesis by its RNA N-glycosidase activity. Ricin toxin, derived from the castor bean *Ricinus communis*, is a prototypic A-B toxin in which the B chain binds to the target cell, and the A chain (RTA) mediates the toxic activity. Ricin B chain (RTB) is a lectin that is responsible for cell agglutination and binds to β -D-galactopyranoside moieties found at the cell surface (e.g., on glycoproteins), allowing the A chain to enter the cell. In turn, the A chain functions enzymatically as an RNA N-glycosidase that depurinates adenine 4324 in the 28S rRNA of the 60S ribosomal subunit. The crystal structure of ricin has been defined.

REFERENCES

- Sharma, S., Podder, S.K. and Karande, A.A. 1999. Comparative studies on kinetics of inhibition of protein synthesis in intact cells by Ricin and a conjugate of Ricin B chain with momordin. *Mol. Cell. Biochem.* 200: 133-141.
- Sandvig, K., van Deurs, B. 2000. Entry of Ricin and Shiga toxin into cells: molecular mechanisms and medical perspectives. *EMBO J.* 19: 5943-5950.
- Candy, L., Peumans, W.J., Menu-Bouaouiche, L., Astoul, C.H., Van Damme, J., Van Damme, E.J., Erard, M. and Rouge, P. 2001. The Gal/GalNAc-specific lectin from the plant pathogenic basidiomycete *Rhizoctonia solani* is a member of the Ricin B family. *iochem. Biophys. Res. Commun.* 282: 655-661.
- Melton-Celsa, A. and O'Brien, A.D. 2003. Plant and bacterial toxins as RNA N-glycosidases. Burns, D.L., Barbieri, J.T., Iglewski, B.H. and Rappuoli, R. eds. *Bacterial Protein Toxins* 245. Am. Soc. Microbiol., Washington, DC.
- Doan, L.G. 2004. Ricin: mechanism of toxicity, clinical manifestations, and vaccine development. A review. *J. Toxicol. Clin. Toxicol.* 42: 201-208.
- Maddaloni, M., Cooke, C., Wilkinson, R., Stout, A.V., Eng, L. and Pincus, S.H. 2004. Immunological characteristics associated with the protective efficacy of antibodies to Ricin. *J. Immunol.* 172: 6221-6228.
- Wu, Y.H., Shih, S.F. and Lin, J.Y. 2004. Ricin triggers apoptotic morphological changes through caspase-3 cleavage of BAT3. *J. Biol. Chem.* 279: 19264-19275.
- McGuinness, C.R. and Mantis, N.J. 2006. Characterization of a novel high-affinity monoclonal immunoglobulin G antibody against the Ricin B subunit. *Infect. Immun.* 74: 3463-3470.
- SWISS-PROT/TrEMBL (P02879). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>.

SOURCE

Ricin B (RB127) is a mouse monoclonal antibody raised against agglutinin of *Ricinus communis* origin.

PRODUCT

Each vial contains 100 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Ricin B (RB127) is recommended for detection of RCA120 and the B-chain fo RCA60 of *Ricinus communis* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

Molecular Weight of Ricin B: 32/120 kDa.

SELECT PRODUCT CITATIONS

- Carter, J.E., Odumosu, O. and Langridge, W.H. 2010. Expression of a ricin toxin B subunit: Insulin fusion protein in edible plant tissues. *Mol. Biotechnol.* 44: 90-100.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.