

FAS (ICO-160): sc-52396

BACKGROUND

Cytotoxic T lymphocyte (CTL)-mediated cytotoxicity constitutes an important component of specific effector mechanisms in immuno-surveillance against virus-infected or transformed cells. Two mechanisms appear to account for this activity, one of which is the perforin-based process. Independently, a FAS-based mechanism involves the transducing molecule FAS (APO-1) and its ligand (FAS-L). The human FAS protein is a cell surface glycoprotein that belongs to a family of receptors that includes CD40, nerve growth factor receptors and tumor necrosis factor receptors. The FAS antigen is expressed on a broad range of lymphoid cell lines, certain ones of which undergo apoptosis in response to treatment with antibody to FAS. These findings strongly imply that targeted cell death is potentially mediated by the intercellular interactions of FAS with its ligand or effectors, and may be critically involved in CTL-mediated cytotoxicity.

REFERENCES

- Henkart, P.A. 1985. Mechanism of lymphocyte-mediated cytotoxicity. *Annu. Rev. Immunol.* 3: 31-58.
- Young, J., Liu, C., Persechini, P. and Cohn, Z. 1988. Perforin-dependent and -independent pathways of cytotoxicity mediated by lymphocytes. *Immunol. Rev.* 103: 161-202.
- Yagita, H., Nakata, M., Kawasaki, A., Shinkai, Y. and Okumura, K. 1992. Role of perforin in lymphocyte-mediated cytolysis. *Adv. Immunol.* 51: 215-242.
- Drappa, J., Brot, N. and Elkon, K. 1993. The FAS protein is expressed at high levels on CD4⁺ CD8⁺ thymocytes and activated mature lymphocytes in normal mice but not in the lupus-prone strain, MRL lpr/lpr. *Proc. Natl. Acad. Sci. USA* 90: 10340-10344.
- Suda, T., Takahashi, T., Golstein, P. and Nagata, S. 1993. Molecular cloning and expression of the FAS ligand, a novel member of the tumor necrosis factor family. *Cell* 75: 1169-1178.
- Kägi, D., Vignaux, F., Ledermann, B., Burki, K., Depraetere, V., Nagata, S., Hengartner, H. and Golstein, P. 1994. FAS and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. *Science* 265: 528-530.
- Hanabuchi, S., Koyanagi, M., Kawasaki, A., Shinohara, N., Matsuzawa, A., Nishimura, Y., Kobayashi, Y., Yonehara, S., Yagita, H. and Okumura, K. 1994. FAS and its ligand in a general mechanism of T cell-mediated cytotoxicity. *Proc. Natl. Acad. Sci. USA* 91: 4930-4934.

CHROMOSOMAL LOCATION

Genetic locus: FAS (human) mapping to 10q24.1; Fas (mouse) mapping to 19 C1.

SOURCE

FAS (ICO-160) is a mouse monoclonal antibody raised against human FAS.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as fluorescein conjugate for flow cytometry, sc-52396 FITC, 100 tests.


APPLICATIONS

FAS (ICO-160) is recommended for detection of FAS of human origin by flow cytometry (1 µg per 1 x 10⁶ cells).

Suitable for use as control antibody for FAS siRNA (h): sc-29311 and FAS siRNA (h2): sc-44260.

Molecular Weight of FAS: 48 kDa.

DATA



FAS (ICO-160): sc-52396. Indirect FCM analysis of HuT 78 cells stained with FAS (ICO-160), followed by PE-conjugated goat anti-mouse IgG: sc-3738. Black line histogram represents the isotype control, normal mouse IgG_{2a}: sc-3878.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.