

CMV pp65 (1-L-11): sc-52401

BACKGROUND

Cytomegalovirus (CMV) is a member of the herpes virus group which includes herpes simplex virus types 1 and 2; Varicella zoster virus, which causes chicken pox; and Epstein Barr virus, which causes infectious mononucleosis. These viruses remain dormant within the body over a long period. In humans, CMV is known as HCMV or human herpesvirus 5 (HHV-5). HHV-5 causes only a brief mononucleosis-like malaise in immunocompetent adults, but may cause severe illness or death in immunosuppressed individuals. CMV pp65 is the early-late lower matrix phosphoprotein of CMV that may be relevant to the etiopathogenesis of scleroderma. CMV pp65 is a major constituent of the CMV virion bodies and is abundantly synthesized during lytic infection. In addition, the CMV pp65 protein is a frequent target for the exceptionally strong CMV-specific CD8⁺ T cell response.

REFERENCES

1. Margraf, S., et al. 2001. Antisense oligonucleotide ISIS 2922 targets IE-expression and prevents HCMV suppression of TSP-1 and TSP-2 expression. *Nucleosides Nucleotides Nucleic Acids* 20: 1425-1428.
2. Khan, N., et al. 2002. Comparative analysis of CD8⁺ T cell responses against human cytomegalovirus proteins pp65 and immediate early 1 shows similarities in precursor frequency, oligoclonality, and phenotype. *J. Infect. Dis.* 185: 1025-1034.
3. Naar, S.P., et al. 2002. Kinetics of HCMV immediate early mRNA expression in stably transfected fibroblasts. *J. Cell Sci.* 115: 321-328.
4. Vlasák, J., et al. 2003. Comparison of hCMV immediate early and CaMV 35S promoters in both plant and human cells. *J. Biotechnol.* 103: 197-202.
5. McGregor, A., et al. 2004. Molecular, biological, and *in vivo* (CMV) homologs of the human CMV matrix proteins pp71 (UL82) and pp65 (UL83). *J. Virol.* 78: 9872-9889.
6. Moss, P. and Khan, N. 2004. CD8⁺ T cell immunity to cytomegalovirus. *Hum. Immunol.* 65: 456-464.
7. Carlsson, B., et al. 2005. Simultaneous generation of cytomegalovirus-specific CD8⁺ and CD4⁺ T lymphocytes by use of dendritic cells comodified with pp65 mRNA and pp65 protein. *J. Infect. Dis.* 192: 1912-1920.
8. Namboodiri, A.M., et al. 2006. Antibodies to human cytomegalovirus protein UL83 in systemic sclerosis. *Clin. Exp. Rheumatol.* 24: 176-178.

SOURCE

CMV pp65 (1-L-11) is a mouse monoclonal antibody raised against cells infected with CMV.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CMV pp65 (1-L-11) is recommended for detection of pp65 of CMV origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of CMV pp65: 65 kDa.

SELECT PRODUCT CITATIONS

1. Arcangeletti, M.C., et al. 2011. Cell-cycle-dependent localization of human cytomegalovirus UL83 phosphoprotein in the nucleolus and modulation of viral gene expression in human embryo fibroblasts *in vitro*. *J. Cell. Biochem.* 112: 307-317.
2. Yu, Y., et al. 2012. Human cytomegalovirus infection induces adipocyte-like lipogenesis through activation of sterol regulatory element binding protein 1. *J. Virol.* 86: 2942-2949.
3. Chuang, J.Y., et al. 2012. Cyr61 increases matrix metalloproteinase-3 expression and cell motility in human oral squamous cell carcinoma cells. *J. Cell. Biochem.* 113: 1977-1986.
4. Lepiller, Q., et al. 2013. HCMV activates the IL-6-JAK-STAT3 axis in Hep G2 cells and primary human hepatocytes. *PLoS ONE* 8: e59591.
5. Yu, Y., et al. 2013. PKR-like endoplasmic reticulum kinase is necessary for lipogenic activation during HCMV infection. *PLoS Pathog.* 9: e1003266.
6. van Domselaar, R., et al. 2013. Granzyme M targets host cell hnRNP K that is essential for human cytomegalovirus replication. *Cell Death Differ.* 20: 419-429.
7. Su, X., et al. 2015. Expression of FABP4, adipin and adiponectin in Paneth cells is modulated by gut *Lactobacillus*. *Sci. Rep.* 5: 18588.
8. Jiang, F., et al. 2015. L-carnitine ameliorates cancer cachexia in mice partly via the carnitine palmitoyltransferase-associated PPAR-γ signaling pathway. *Oncol. Res. Treat.* 38: 511-516.
9. Zwezdaryk, K.J., et al. 2015. Human cytomegalovirus infection of human adipose-derived stromal/stem cells restricts differentiation along the adipogenic lineage. *Adipocyte* 5: 53-64.
10. Caruana, I., et al. 2015. K562-derived whole-cell vaccine enhances antitumor responses of CAR-redirection virus-specific cytotoxic T lymphocytes *in vivo*. *Clin. Cancer Res.* 21: 2952-2962.
11. Vysochan, A., et al. 2017. ACS2-mediated acetyl-CoA synthesis from acetate is necessary for human cytomegalovirus infection. *Proc. Natl. Acad. Sci. USA* 114: E1528-E1535.
12. Yan, C., et al. 2018. Smad ubiquitination regulatory factor 1 (Smurf1) promotes thyroid cancer cell proliferation and migration via ubiquitin-dependent degradation of Kisspeptin-1. *Cell. Physiol. Biochem.* 49: 2047-2059.
13. Yu, J.T., et al. 2021. DNA methylation of FTO promotes renal inflammation by enhancing m⁶A of PPAR-α in alcohol-induced kidney injury. *Pharmacol. Res.* 163: 105286.

RESEARCH USE

For research use only, not for use in diagnostic procedures.