

IL-10 (JES5-16E3): sc-52560

BACKGROUND

Interleukin-10, or IL-10, is a 178 amino acid protein that is primarily secreted by TH2 clones. IL-10 has dual functions, the first of which is the suppression of cytokine production by TH1 clones responding to antigen presented by monocyte and macrophage antigen presenting cells (APCs). The second function consists of the inhibition of response of cytokine targeted cells, possibly by the downregulation of CD25 (the IL-2 receptor) on macrophages and B lymphocytes. Human and murine IL-10 exhibit 81% sequence identity at the amino acid level and share 73% identity at the cDNA level. Both human and murine IL-10 are acid-labile and exist as non-covalently-linked homodimers in solution. IL-10 exerts its biological activity through the IL-10 receptor (IL-10R), a glycoprotein whose expression can be induced in cultured macrophages and fibroblasts by lipopolysaccharide (LPS) stimulation. IL-10 expression has been shown to be elevated in HIV-1 infected individuals and has been implicated in the progression of the disease.

REFERENCES

- Feng, L., et al. 1993. Molecular cloning of rat cytokine synthesis inhibitory factor (IL-10) cDNA and expression in spleen and macrophages. *Biochem. Biophys. Res. Commun.* 192: 452-458.
- Cohen, S.B., et al. 1994. IL-10 enhances expression of the IL-2 receptor α chain on T cells. *Immunology* 83: 329-332.
- Weber-Nordt, R.M., et al. 1994. Lipopoly-saccharide-dependent induction of IL-10 receptor expression on murine fibroblasts. *J. Immunol.* 153: 3734-3744.
- Bromberg, J.S. 1995. IL-10 immunosuppression in transplantation. *Curr. Opin. Immunol.* 7: 639-643.
- Finnegan, A., et al. 1996. IL-10 cooperates with TNF- α to activate HIV-1 from latently and acutely infected cells of monocyte/macrophage lineage. *J. Immunol.* 156: 841-851.
- Ludewig, B., et al. 1996. Transmission of HIV-1 from productively infected mature Langerhans cells to primary CD4⁺ T lymphocytes results in altered T cell responses with enhanced production of IFN- γ and IL-10. *Virology* 215: 51-60.
- Fleming, S.D. and Campbell, P.A. 1996. Macrophages have cell surface IL-10 that regulates macrophage bactericidal activity. *J. Immunol.* 156: 1143-1150.
- Min, C.K., et al. 2007. IL-10-transduced bone marrow mesenchymal stem cells can attenuate the severity of acute graft-versus-host disease after experimental allogeneic stem cell transplantation. *Bone Marrow Transplant.* 39: 637-645.

CHROMOSOMAL LOCATION

Genetic locus: Il10 (mouse) mapping to 1 E4.

SOURCE

IL-10 (JES5-16E3) is a rat monoclonal antibody raised against full length IL-10 of mouse origin.

PRODUCT

Each vial contains 100 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


IL-10 (JES5-16E3) is recommended for detection of IL-10 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-10 siRNA (m): sc-39635, IL-10 shRNA Plasmid (m): sc-39635-SH and IL-10 shRNA (m) Lentiviral Particles: sc-39635-V.

Molecular Weight of IL-10 monomer: 20 kDa.

Molecular Weight of IL-10 dimer: 37 kDa.

DATA



IL-10 (JES5-16E3): sc-52560. Western blot analysis of mouse recombinant IL-10 fusion protein.

SELECT PRODUCT CITATIONS

- Yin, Z., et al. 2000. Dominance of IL-12 over IL-4 in $\gamma\delta$ T cell differentiation leads to default production of IFN- γ : failure to down-regulate IL-12 receptor β_2 -chain expression. *J. Immunol.* 164: 3056-3064.
- Rahayu, R.P., et al. 2020. Combinations of propolis and Ca(OH)₂ in dental pulp capping treatment for the stimulation of reparative dentin formation in a rat model. *F1000Res.* 9: 308.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **IL-10 (A-2): sc-365858** for IL-10 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.