

RLA-DR (RDR34): sc-52610

BACKGROUND

Several class II α and β chain genes of the rabbit major histocompatibility complex have been classified into three distinct subregions, R-DP, R-DQ and R-DR, based on their homology to the corresponding HLA-DP, -DQ and -DR genes. Studies indicate that the rabbit germline contains a total of approximately seven class II β genes, one DQ β , one DP β and five DR β . R-DQ and R-DR molecules show expression on cell surfaces, whereas R-DP molecules exhibit low levels of expression in the spleen. The constitutive coexpression of the major histocompatibility complex (MHC) class II genes in B lymphocytes requires positive, *trans*-acting transcriptional factors, although the mechanism by which the *trans*-acting factors exert their effect on gene transcription is unknown.

REFERENCES

1. Sittisombut, N. and Knight, K.L. 1986. Rabbit major histocompatibility complex. I. Isolation and characterization of three subregions of class II genes. *J. Immunol.* 136: 1871-1875.
2. Kulaga, H., Sogn, J.A., Weissman, J.D., Marche, P.N., LeGuern, C., Long, E.O. and Kindt, T.J. 1987. Expression patterns of MHC class II genes in rabbit tissues indicate close homology to human counterparts. *J. Immunol.* 139: 587-592.
3. Sittisombut, N. 1988. Two distinct nuclear factors bind the conserved regulatory sequences of a rabbit major histocompatibility complex class II gene. *Mol. Cell. Biol.* 8: 2034-2041.
4. Sittisombut, N., Tissot, R.G. and Knight, K.L. 1989. Rabbit major histocompatibility complex. III. Multiple class II DR β genes and restriction fragment length polymorphism of the class II α and β genes. *J. Immunogenet.* 16: 63-75.
5. Spieker-Polet, H., Sittisombut, N., Yam, P.C. and K.L. Knight. 1990. Rabbit major histocompatibility complex IV. Expression of major histocompatibility complex class II genes. *J. Immunogenet.* 17: 123-132.
6. Wilkinson, J.M., McDonald, G., Smith, S., Galea-Lauri, J., Lewthwaite, J., Henderson, B. and Revell, P.A. 1993. Immunohistochemical identification of leucocyte populations in normal tissue and inflamed synovium of the rabbit. *J. Pathol.* 170: 315-320.
7. Fain, M.A., Zhao, T. and Kindt, T.J. 2001. Improved typing procedure for the polymorphic single-copy RLA-DQA gene of the rabbit reveals a new allele. *Tissue Antigens* 57: 332-328.
8. Jabrane-Ferrat, N., Nekrep, N., Tosi, G., Esserman, L. and Peterlin, B.M. 2003. DNA-bound activators? *Int. Immunol.* 15: 467-475.

SOURCE

RLA-DR (RDR34) is a mouse monoclonal antibody raised against Leukemic B-cell line of rabbit origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 100 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RLA-DR (RDR34) is available conjugated phycoerythrin (sc-52610 PE, 100 tests in 2 ml), for IF, IHC(P) and FCM.

APPLICATIONS

RLA-DR (RDR34) is recommended for detection of RLA-DR of rabbit origin by immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.