

# Hep C NS3 (12-15): sc-52805

## BACKGROUND

The Hep C (Hepatitis C) is a small, enveloped, single-stranded, positive sense RNA virus belonging to the family *Flaviviridae*. Transmission of the virus occurs when blood from an infected individual enters the body of an uninfected individual. Hep C primarily replicates within hepatocytes in the liver, and circulating Hep C particles bind to receptors on the surface and enter these cells. Hep C replicates quickly, producing approximately one trillion particles each day in infected individuals. Hep C RNA polymerase has no proofreading function, so the virus has an exceptionally high mutation rate which may help it elude the host's immune system. Hep C infection results in chronic infections, liver cirrhosis and hepatocellular carcinoma in most people. Hep C NS3 (nonstructural protein 3) has both protease and helicase activity and is essential for Hep C replication and proliferation.

## REFERENCES

1. Watashi, K. and Shimotohno, K. 2003. The roles of hepatitis C virus proteins in a novel action mechanism of the HCV core protein on gene regulation by nuclear hormone receptors. *Cancer Sci.* 94: 937-943.
2. Acosta-Rivero, N., Rodriguez, A., Musacchio, A., Falcón, V., Suarez, V.M., Chavez, L., Morales-Grillo, J. and Duenas-Carrera, S. 2004. Nucleic acid binding in *Pichia pastoris*. *Biochem. Biophys. Res. Commun.* 323: 926-931.
3. Sansonno, D., Lauletta, G. and Dammacco, F. 2004. Detection and quantitation of HCV core protein in single hepatocytes by means of laser capture microdissection and enzyme-linked immunosorbent assay. *J. Viral Hepat.* 11: 27-32.
4. Umehara, T., Fukuda, K., Nishikawa, F., Sekiya, S., Kohara, M., Hasegawa, T. and Nishikawa, S. 2004. Designing and analysis of a potent bi-functional aptamers that inhibit protease and helicase activities of HCV NS3. *Nucleic Acids Symp. Ser.* 48: 195-196.
5. Alisi, A., Mele, R., Spaziani, A., Tavolaro, S., Palescandolo, E. and Balsano, C. 2005. Thr 446 phosphorylation of PKR by HCV core protein deregulates G<sub>2</sub>/M phase HCC cells. *J. Cell. Physiol.* 205: 25-31.
6. Carabaich, A., Ruvoletto, M., Bernardinello, E., Tono, N., Cavalletto, L., Chemello, L., Gatta, A. and Pontisso, P. 2005. Profiles of HCV core protein and viremia in chronic hepatitis C: possible protective role of core antigen in liver damage. *J. Med. Virol.* 76: 55-60.
7. Gu, J., Wang, L., Che, Y., Liu, L., Jiang, L., Dong, S., Li, W. and Li, Q. 2005. Morphological alteration and biological properties of hepatocytes not related to tumorigenesis following transfection with HCV core protein. *J. Viral Hepat.* 12: 20-26.
8. Kimball, P., Verbeke, S. and Shiffman, M. 2005. HCV core protein augments cyclosporine immunosuppression. *Transplant Proc.* 37: 652-653.
9. Alvarez-Lajonchere, L., González, M., Alvarez-Obregón, J.C., Guerra, I., Viña, A., Acosta-Rivero, N., Musacchio, A., Morales, J. and Dueñas-Carrera, S. 2006. Hepatitis C virus (HCV) core protein enhances the immunogenicity of a co-delivered DNA vaccine encoding HCV structural antigens in mice. *Biotechnol. Appl. Biochem.* 44: 9-17.

## SOURCE

Hep C NS3 (12-15) is a mouse monoclonal antibody raised against a chimeric polyprotein corresponding to at least 60 amino acids of the NS3 region of Chinese Hep C strains.

## PRODUCT

Each vial contains 100 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Hep C NS3 (12-15) is recommended for detection of an epitope corresponding to amino acids 1378-1458 of the NS3 region of Hep C origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000); non cross-reactive with core region or other non-structural regions.

Molecular Weight of Hep C NS3: 70 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.