

SERCA1 (IIH11): sc-53011

BACKGROUND

ATP dependent calcium pumps are responsible, in part, for the maintenance of low cytoplasmic free calcium concentrations. The ATP pumps that reside in intracellular organelles are encoded by a family of structurally related enzymes, termed the sarcoplasmic or endoplasmic reticulum calcium (SERCA) ATPases. The sarcoplasmic reticulum of striated muscle is a specialized intracellular membrane system that plays a critical role in the contraction and relaxation of muscle. The SERCAs mediate Ca^{2+} uptake into intracellular stores. SERCA-mediated Ca^{2+} uptake induces and maintains muscular relaxation. The SERCA1 gene is exclusively expressed in type II (fast) skeletal muscle. The SERCA2 gene is subject to tissue-dependent processing which is responsible for the generation of the SERCA2a muscle-specific form expressed in type I (slow) skeletal, cardiac and smooth muscle, and the SERCA2b isoform expressed in all cell types. The SERCA3 gene is not as well characterized and is found in non-muscle cells. SERCA2 plays an important part in regulating cardiac contractile function. SERCA3 is an isoform expressed in several cell types including platelets, lymphoid cells and mast cells. SERCA1, SERCA2 and SERCA3 all undergo alternative splicing.

REFERENCES

1. Poch, E., et al. 1998. Functional characterization of alternatively spliced human SERCA3 transcripts. *Am. J. Physiol., Cell Physiol.* 275: C1449-C1458.
2. Anger, M., et al. 1998. Cellular distribution of Ca^{2+} pumps and Ca^{2+} release channels in rat cardiac hypertrophy induced by aortic stenosis. *Circulation* 98: 2477-2486.
3. Loukianov, E., et al. 1998. Enhanced myocardial contractility and increased Ca^{2+} transport function in transgenic hearts expressing the fast-twitch skeletal muscle sarcoplasmic reticulum Ca^{2+} -ATPase. *Circ. Res.* 83: 889-897.
4. Aubier, M. and Viires, N. 1998. Calcium ATPase and respiratory muscle function. *Eur. Respir. J.* 11: 758-766.
5. Bobe, R., et al. 1998. Expression of two isoforms of the third sarco/endoplasmic reticulum Ca^{2+} ATPase (SERCA3) in platelets. Possible recognition of the SERCA3b isoform by the PL/IM430 monoclonal antibody. *FEBS Lett.* 423: 259-264.

CHROMOSOMAL LOCATION

Genetic locus: *Atp2a1* (mouse) mapping to 7 F3.

SOURCE

SERCA1 (IIH11) is a mouse monoclonal antibody raised against purified skeletal muscle sarcoplasmic reticulum of rabbit origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SERCA1 (IIH11) is recommended for detection of SERCA1 of mouse, rat, rabbit, canine and guinea pig origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for SERCA1 siRNA (m): sc-36483, SERCA1 shRNA Plasmid (m): sc-36483-SH and SERCA1 shRNA (m) Lentiviral Particles: sc-36483-V.


Molecular Weight of SERCA1: 110 kDa.

Positive Controls: A-10 cell lysate: sc-3806, rat skeletal muscle extract: sc-364810 or mouse skeletal muscle extract: sc-364250.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



SERCA1 (IIH11): sc-53011. Western blot analysis of SERCA1 expression in rat skeletal muscle tissue extract.



SERCA1 (IIH11): sc-53011. Western blot analysis of SERCA1 expression in mouse skeletal muscle tissue extract.

SELECT PRODUCT CITATIONS

1. Strosova, M., et al. 2009. Modulation of SERCA in the chronic phase of adjuvant arthritis as a possible adaptation mechanism of redox imbalance. *Free Radic. Res.* 43: 852-864.
2. Brulé, C., et al. 2010. Proteomic study of calpain interacting proteins during skeletal muscle aging. *Biochimie* 92: 1923-1933.
3. Strosova, M.K., et al. 2011. Modulation of sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase activity and oxidative modification during the development of adjuvant arthritis. *Arch. Biochem. Biophys.* 511: 40-47.

RESEARCH USE

For research use only, not for use in diagnostic procedures.