SANTA CRUZ BIOTECHNOLOGY, INC.

Ku86 (5D5.8): sc-53021



BACKGROUND

The Ku protein is localized in the nucleus and is composed of subunits referred to as Ku70 (p70) and Ku86 (p86) which is also known by the synonym Ku80 or (p80). Ku was first described as an autoantigen to which antibodies were produced in a patient with scleroderma polymyositis overlap syndrome, and was later found in the sera of patients with other rheumatic diseases. Both subunits of the Ku protein have been cloned, and a number of functions have been proposed for Ku, including cell signaling, DNA replication and transcriptional activation. Ku is involved in Pol II-directed transcription by virtue of its DNA binding activity, serving as the regulatory component of the DNA-associated protein kinase that phosphorylates Pol II and transcription factor Sp. Ku proteins also activate transcription from the U1 small nuclear RNA and the human transferrin receptor gene promoters. A Ku-related protein designated the enhancer 1 binding factor (E1BF), composed of two subunits, has been identified as a positive regulator of RNA polymerase I transcription initiation.

REFERENCES

- Mimori, T., et al. 1981. Characterization of a high molecular weight acidic nuclear protein recognized by autoantibodies in sera from patients with polymyositis-scleroderma overlap. J. Clin. Invest. 68: 611-620.
- Mimori, T., et al. 1986. Characterization of the DNA-binding protein antigen Ku recognized by autoantibodies from patients with rheumatic disorders. J. Biol. Chem. 261: 2274-2278.
- Chan, J.Y.C., et al. 1989. Cloning and characterization of a cDNA that encodes a 70-kDa novel human thyroid autoantigen. J. Biol. Chem. 264: 3651-3654.
- 4. Reeves, W.H., et al. 1989. Molecular cloning of cDNA encoding the p70 (Ku) lupus autoantigen. J. Biol. Chem. 264: 5047-5052.
- 5. Yaneva, M., et al. 1989. cDNA-derived amino acid sequence of the 86-kDa subunit of the Ku antigen. J. Biol. Chem. 264: 13407-13411.
- Prabhakar, B.S., et al. 1990. Cell surface expression of the 70-kD component of Ku, a DNA-binding nuclear antigen. J. Clin. Invest. 86: 1301-1305.
- 7. Stuiver, M.H, et al. 1990. The autoantigen Ku is indistinguishable from NF IV, a protein forming multimeric protein-DNA complexes. J. Exp. Med. 172: 1049-1054.
- Gottlieb, T.M., et al. 1993. The DNA-dependent protein kinases: requirement for DNA ends and association with Ku antigen. Cell 72: 131-142.
- Hoff, C.M., et al. 1994. Enhancer 1 binding factor, a Ku-related protein, is a positive regulator of RNA polymerase I transcription initiation. Proc. Natl. Acad. Sci. USA 91: 762-766.

CHROMOSOMAL LOCATION

Genetic locus: XRCC5 (human) mapping to 2q35.

SOURCE

Ku86 (5D5.8) is a mouse monoclonal antibody raised against HeLa cell extracts.

PRODUCT

Each vial contains 200 μg lgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Ku86 (5D5.8) is recommended for detection of Ku86 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Ku86 siRNA (h): sc-29384, Ku86 shRNA Plasmid (h): sc-29384-SH and Ku86 shRNA (h) Lentiviral Particles: sc-29384-V.

Molecular Weight of Ku86: 86 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, C32 whole cell lysate: sc-2205 or A-431 whole cell lysate: sc-2201.

DATA





Ku86 (5D5.8): sc-53021. Western blot analysis of Ku86 expression in A-431 (A), HeLa (B) and C32 (C) whole cell lysates and A-431 (D), HeLa (E) and C32 (F) nuclear extracts.

Ku86 (5D5.8): sc-53021. Western blot analysis of Ku86 expression in Hep G2 (**A**) and Jurkat (**B**) nuclear extracts.

SELECT PRODUCT CITATIONS

 Yin, H., et al. 2020. UBE2T promotes radiation resistance in non-small cell lung cancer via inducing epithelial-mesenchymal transition and the ubiguitination-mediated FOXO1 degradation. Cancer Lett. 494: 121-131.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **Ku86 (B-1): sc-5280** for Ku86 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.