

Ly-49F (HBF-719): sc-53023

BACKGROUND

The activity of natural killer (NK) cells is regulated by members of multiple receptor families that recognize class I MHC molecules, such as the killer cell inhibitory receptor/leukocyte immunoglobulin-like receptor (KIR/LIR) family and the C-type lectin superfamily. The KIR/LIR family includes p91A (also designated pp130 or PIR-B, for paired Immunoglobulin-like receptor-B) and p91B (also designated PIR-A). p91A acts as an inhibitory receptor through interactions with SHP-1, whereas p91B acts as an activating receptor. CD94, NKG2 and Ly-49 are members of the C-type lectin superfamily of type II membrane glycoproteins. CD94 forms heterodimers with NKG2 isoforms on the surface of NK cells, whereas Ly-49 isoforms form homodimers. There are at least 21 Ly-49 proteins, ranging from 230 to 280 amino acids in length. The Ly-49 isoforms, which are clustered on human chromosome 16, all function to present the C-type lectin domain on the cell membrane. NKG2-D, expressed on NK cells, $\gamma\delta$ T cells, and CD8⁺ $\alpha\beta$ T cells, is a receptor for the stress inducible protein MICA, an antigen frequently expressed in epithelial tumors.

REFERENCES

1. Ljunggren, H.G. and Kärre, K. 1990. In search of the "missing self": MHC molecules and NK cell recognition. *Immunol. Today* 11: 237-244.
2. Ortaldo, J.R., Winkler-Pickett, R., Mason, A.T. and Mason, L.H. 1998. The Ly-49 family: regulation of cytotoxicity and cytokine production in murine CD3⁺ cells. *J. Immunol.* 160: 1158-1165.
3. Robson MacDonald, H., Lees, R.K. and Held, W. 1998. Developmentally regulated ext and selection of NK1.1⁺ T cells. *J. Exp. Med.* 187: 2109-2114.
4. Manilay, J.O., Wanek, G.L. and Sykes, M. 1999. Levels of Ly-49 receptor expression with MHC ligands: evidence against receptor calibration to a "useful" level. *J. Immunol.* 163: 2628-2633.
5. Naper, C., Ryan, J.C., Kirsch, R., Butcher, G.W., Rolstad, B., and Vaage, J.T. 1999. Genes in two major histocompatibility complex class I regions control selection, phenotype, and function of a rat Ly-49 natural killer cell subset. *Eur. J. Immunol.* 29: 2046-2053.
6. Coles, M.C., McMahon, C.W., Takizawa, H. and Raulet, D.H. 2000. Memory CD8 T lymphocytes express inhibitory MHC-specific Ly-49 receptors. *Eur. J. Immunol.* 30: 236-244.
7. Kane, K.P., Silver, E.T. and Hazes, B. 2001. Specificity and function of activating Ly-49 receptors. *Immunol. Rev.* 181: 104-114.
8. Kunz, B. and Held, W. 2001. Positive and negative roles of the *trans*-acting T ce of distinct Ly-49 MHC class I receptors by NK cells. *J. Immunol.* 166: 6181-6187.
9. Kane, K.P., Lavender, K.J. and Ma, B.J. 2005. Ly-49 receptors and their functions. *Crit. Rev. Immunol.* 24: 321-348.

CHROMOSOMAL LOCATION

Genetic locus: Klra6 (mouse) mapping to 6 F3.

SOURCE

Ly-49F (HBF-719) is a mouse monoclonal antibody raised against the B6 allele of Ly-49F of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Ly-49F (HBF-719) is available conjugated to either phycoerythrin (sc-53023 PE) or fluorescein (sc-53023 FITC), 200 μ g/ml, for IF, IHC(P) and FCM.

APPLICATIONS

Ly-49F (HBF-719) is recommended for detection of Ly-49F of mouse origin by flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for Ly-49F siRNA (m): sc-72051, Ly-49F shRNA Plasmid (m): sc-72051-SH and Ly-49F shRNA (m) Lentiviral Particles: sc-72051-V.

Molecular Weight of Ly-49F: 31 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.