# SANTA CRUZ BIOTECHNOLOGY, INC.

# His-Tag (AD1.1.10): sc-53073



## BACKGROUND

Plasmid vectors for the expression of coding regions of eukaryotic genes in bacterial, insect and mammalian hosts are in common usage; such expression vectors are frequently used to encode hybrid fusion proteins consisting of a eukaryotic target protein and a specialized region designed to aid in the purification and visualization of the target protein. A system that has proven to be very successful relies on the insertion of a six histidine (His6) sequence in the N-terminus of the encoded protein, allowing for efficient coupling to Ni<sup>2+</sup>-chelating resins and purification by single step affinity chromatography. This polyhistidine sequence can then be removed by specific cleavage at sites recognized by enzymes such as thrombin or enterokinase, permitting the separation of the target protein from the polyhistidine tag. Visualization of such fusion proteins can be achieved by utilizing antibodies generated against specific peptide sequences downstream from the multiple cloning site.

### SOURCE

His-Tag (AD1.1.10) is a mouse monoclonal antibody raised against a 6x His-tagged polypeptide.

### PRODUCT

Each vial contains 200  $\mu$ g lgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

His-Tag (AD1.1.10) is available conjugated to agarose (sc-53073 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-53073 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-53073 PE), fluorescein (sc-53073 FITC), Alexa Fluor<sup>®</sup> 488 (sc-53073 AF488), Alexa Fluor<sup>®</sup> 546 (sc-53073 AF546), Alexa Fluor<sup>®</sup> 594 (sc-53073 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-53073 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-53073 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-53073 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

#### **APPLICATIONS**

His-Tag (AD1.1.10) is recommended for detection of fusion proteins encoded by polyhistidine expression vectors by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

# **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# DATA



His-Tag (AD1.1.10): sc-53073. Western blot analysis of His-tagged fusion proteins (**A**,**B**).

#### SELECT PRODUCT CITATIONS

- Zhang, X., et al. 2008. Histone deacetylase-related protein inhibits AESmediated neuronal cell death by direct interaction. J. Neurosci. Res. 86: 2423-2431.
- Liu, Y.C., et al. 2011. The role of small ubiquitin-like modifier-interacting motif in the assembly and regulation of metal-responsive transcription factor 1. J. Biol. Chem. 286: 42818-42829.
- Mao, Y., et al. 2012. Subunit organization of the membrane-bound HIV-1 envelope glycoprotein trimer. Nat. Struct. Mol. Biol. 19: 893-899.
- 4. Alvarez, L.A., et al. 2013. Cj1411c encodes for a cytochrome P450 involved in *Campylobacter jejuni* 81-176 pathogenicity. PLoS ONE 8: e75534.
- 5. Chae, Y.C., et al. 2014. Inhibition of FoxO1 acetylation by INHAT subunit SET/TAF-I $\beta$  induces p21 transcription. FEBS Lett. 588: 2867-2873.
- 6. Zhang, S., et al. 2015. PCAF-mediated Akt1 acetylation enhances the proliferation of human glioblastoma cells. Tumour Biol. 36: 1455-1462.
- 7. Avila, E.E., et al. 2016. An *Entamoeba histolytica* ADP-ribosyl transferase from the diphtheria toxin family modifies the bacterial elongation factor Tu. Mol. Biochem. Parasitol. 207: 68-74.
- 8. Gao, J., et al. 2017. Kindlin supports platelet integrin αllbβ3 activation by interacting with paxillin. J. Cell Sci. 130: 3764-3775.
- Kim, Y.R., et al. 2018. *Toxoplasma gondii* GRA8 induces ATP5A1-SIRT3mediated mitochondrial metabolic resuscitation: a potential therapy for sepsis. Exp. Mol. Med. 50: e464.
- Men, Y., et al. 2019. Exosome reporter mice reveal the involvement of exosomes in mediating neuron to astroglia communication in the CNS. Nat. Commun. 10: 4136.
- Kim, J.S., et al. 2020. Toxoplasma gondii GRA8-derived peptide immunotherapy improves tumor targeting of colorectal cancer. Oncotarget 11: 62-73.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.