

IgD (OX60): sc-53082

BACKGROUND

Immunoglobulins are four-chain, Y-shaped, monomeric structures comprised of two identical heavy chains and two identical light chains held together through interchain disulfide bonds. The chains form two domains, the Fab (antigen binding) fragment and the Fc (constant) fragment. Immunoglobulin D (IgD) exists as a monomer with δ heavy chains and either κ or λ light chains. It plays a biological role as a transmembrane receptor molecule, co-expressed with IgM on the surface of mature/naive B cells. In particular, it is found on spleen B cell surfaces. Compared to IgM, IgD exists in much lower numbers and is not expressed on immature B cells. IgD surface expression on B cells is regulated in part by IL-27. In mice, the inhibition of this immunoglobulin isotype does not cause a significant change to the immune system.

REFERENCES

1. Takayasu, T., et al. 1980. Amino acid sequence and location of the three glycopeptides in the Fc region of human immunoglobulin D. *Biochem. Biophys. Res. Commun.* 97: 635-641.
2. Lin, L.C. and Putnam, F.W. 1981. Primary structure of the Fc region of human immunoglobulin D: implications for evolutionary origin and biological function. *Proc. Natl. Acad. Sci. USA* 78: 504-508.
3. Shinoda, T., et al. 1981. Complete amino acid sequence of the Fc region of a human δ chain. *Proc. Natl. Acad. Sci. USA* 78: 785-789.
4. Putnam, F.W., et al. 1981. Amino acid sequence of the first constant region domain and the hinge region of the δ heavy chain of human IgD. *Proc. Natl. Acad. Sci. USA* 78: 6168-6172.
5. Takayasu, T., et al. 1982. Amino acid sequence of galactosamine-containing glycopeptides in the hinge region of a human immunoglobulin D. *Biochem. Biophys. Res. Commun.* 105: 1066-1071.
6. Naiem, M., et al. 1982. The value of immunohistological screening in the production of monoclonal antibodies. *J. Immunol. Methods* 50: 145-160.
7. Ohta, Y. and Flajnik, M. 2006. IgD, like IgM, is a primordial immunoglobulin class perpetuated in most jawed vertebrates. *Proc. Natl. Acad. Sci. USA* 103: 10723-10728.
8. Zhao, Y., et al. 2006. Identification of IgF, a hinge-region-containing Ig class, and IgD in *Xenopus tropicalis*. *Proc. Natl. Acad. Sci. USA* 103: 12087-12092.
9. Boumendjel, A., et al. 2006. IL-27 induces the production of IgG1 by human B cells. *Eur. Cytokine Netw.* 17: 281-289.

CHROMOSOMAL LOCATION

Genetic locus: Ighd (mouse) mapping to 12 F1.

SOURCE

IgD (OX60) is a mouse monoclonal antibody raised against purified IgD of rat origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IgD (OX60) is available conjugated to agarose (sc-53082 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-53082 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-53082 PE), fluorescein (sc-53082 FITC), Alexa Fluor® 488 (sc-53082 AF488), Alexa Fluor® 546 (sc-53082 AF546), Alexa Fluor® 594 (sc-53082 AF594) or Alexa Fluor® 647 (sc-53082 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-53082 AF680) or Alexa Fluor® 790 (sc-53082 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

IgD (OX60) is recommended for detection of IgD Chain C of mouse origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 \times 10⁶ cells).

Molecular Weight of IgD heavy (δ) chain: 44-80 kDa.


Molecular Weight of IgD light (κ/λ) chain: 21-25 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 2) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



IgD (OX60): sc-53082. Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing membrane and cytoplasmic staining of subset of cells in white pulp and cells in red pulp (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing membrane and cytoplasmic staining of cells in germinal center and cells in non-germinal center (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.