

Integrin $\alpha 3$ (F35 177-1): sc-53354

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, Collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as Fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis. The Integrin $\alpha 3$ chain, also known as very late (activation) antigen 3 (VLA-3), very common antigen 2 (VCA-2), extracellular matrix receptor 1 (ECMR1) and galactoprotein b3 (GAPB3), undergoes posttranslational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains that join with $\beta 1$ to form an integrin that interacts with many extracellular-matrix proteins.

REFERENCES

1. Tsuji, T., et al. 1991. Identification of human galactoprotein b3, an oncogenic transformation-induced membrane glycoprotein, as VLA-3 α subunit: the primary structure of human Integrin $\alpha 3$. *J. Biochem.* 109: 659-665.
2. Hynes, R.O. 1992. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 69: 11-25.
3. Berdichevsky, F., et al. 1994. Branching morphogenesis of human mammary epithelial cells in collagen gels. *J. Cell Sci.* 107: 3557-3568.
4. Miyamoto, S., et al. 1995. Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. *Science* 267: 883-885.
5. Clark, E.A., et al. 1995. Integrins and signal transduction pathways: the road taken. *Science* 268: 233-239.
6. Sheppard, D. 1996. Epithelial integrins. *Bioessays* 18: 655-660.
7. Juliano, R. 1996. Cooperation between soluble factors and integrin-mediated cell anchorage in the control of cell growth and differentiation. *Bioessays* 18: 911-917.
8. de Melker, A.A., et al. 1997. The A and B variants of the $\alpha 3$ Integrin subunit: tissue distribution and functional characterization. *Lab. Invest.* 76: 547-563.
9. Hirotsaki, T., et al. 2000. Structural requirement of carboxyl-terminal globular domains of Laminin $\alpha 3$ chain for promotion of rapid cell adhesion and migration by Laminin-5. *J. Biol. Chem* 275: 22495-22502.

CHROMOSOMAL LOCATION

Genetic locus: ITGA3 (human) mapping to 17q21.33.

SOURCE

Integrin $\alpha 3$ (F35 177-1) is a mouse monoclonal antibody raised against MTSV1-7 HYE-breast epithelial cells of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Integrin $\alpha 3$ (F35 177-1) is recommended for detection of Integrin $\alpha 3$ of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Integrin $\alpha 3$ siRNA (h): sc-35684, Integrin $\alpha 3$ shRNA Plasmid (h): sc-35684-SH and Integrin $\alpha 3$ shRNA (h) Lentiviral Particles: sc-35684-V.

Molecular Weight of Integrin $\alpha 3$: 150 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Integrin $\alpha 3$ (A-3): sc-374242** for Integrin $\alpha 3$ antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.