

Integrin $\alpha 2$ (P1E6): sc-53502

BACKGROUND

Integrins are heterodimers composed of non-covalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as fibrinogen, or to counter-receptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis. Integrin $\alpha 2$ is responsible for adhesion of platelets and other cells to collagens. Modulation of collagen and collagenase gene expression force generation and organization of newly synthesized extracellular matrix.

CHROMOSOMAL LOCATION

Genetic locus: ITGA2 (human) mapping to 5q11.2.

SOURCE

Integrin $\alpha 2$ (P1E6) is a mouse monoclonal antibody raised against HT-1080 fibrosarcoma cells of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Integrin $\alpha 2$ (P1E6) is recommended for detection of Integrin $\alpha 2$ of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Integrin $\alpha 2$ siRNA (h): sc-29371, Integrin $\alpha 2$ shRNA Plasmid (h): sc-29371-SH and Integrin $\alpha 2$ shRNA (h) Lentiviral Particles: sc-29371-V.


Molecular Weight of Integrin $\alpha 2$: 150 kDa.

Positive Controls: human platelet extract: sc-363773, CCRF-CEM cell lysate: sc-2225 or CCRF-HSB-2 cell lysate: sc-2265.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Integrin $\alpha 2$ (P1E6): sc-53502. Western blot analysis of Integrin $\alpha 2$ expression in human platelet extract (A) and human PBL whole cell lysate (B) under non-reducing conditions.

SELECT PRODUCT CITATIONS

- Vincourt, J.B., et al. 2010. C-propeptides of procollagens I α 1 and II that differentially accumulate in enchondromas versus chondrosarcomas regulate tumor cell survival and migration. *Cancer Res.* 70: 4739-4748.
- Sung, B.H., et al. 2011. Cortactin controls cell motility and lamellipodial dynamics by regulating ECM secretion. *Curr. Biol.* 21: 1460-1469.
- Sato-Nishiuchi, R., et al. 2012. Polydom/SVEP1 is a ligand for Integrin $\alpha 9\beta 1$. *J. Biol. Chem.* 287: 25615-25630.
- Bartolomé, R.A., et al. 2014. Cadherin-17 interacts with $\alpha 2\beta 1$ Integrin to regulate cell proliferation and adhesion in colorectal cancer cells causing liver metastasis. *Oncogene* 33: 1658-1669.
- Hozumi, K., et al. 2015. Suppression of cell adhesion through specific integrin crosstalk on mixed peptide-polysaccharide matrices. *Biomaterials* 37: 73-81.
- Bartolomé, R.A., et al. 2016. VE-cadherin RGD motifs promote metastasis and constitute a potential therapeutic target in melanoma and breast cancers. *Oncotarget* 8: 215-227.
- Bartolomé, R.A., et al. 2021. CDH6-activated $\alpha 11\beta 3$ crosstalks with $\alpha 2\beta 1$ to trigger cellular adhesion and invasion in metastatic ovarian and renal cancers. *Mol. Oncol.* 15: 1849-1865.
- Fayad, R., et al. 2021. EFA6B regulates a stop signal for collective invasion in breast cancer. *Nat. Commun.* 12: 2198.

RESEARCH USE

For research use only, not for use in diagnostic procedures.