

prefoldin 6 (1D10-4D11): sc-53519

BACKGROUND

Molecular chaperones are proteins that assist in the correct folding of other proteins in the crowded molecular environment that exists in living cells. Within this class of proteins, a key role is played by chaperonins, multisubunit toroidal (i.e. doughnut-shaped) assemblies that undergo major ATP-dependent conformational changes as part of the mechanism of facilitated folding. Prefoldin is a heterohexameric chaperone protein which has the ability to capture unfolded Actin. Six prefoldin polypeptides, prefoldin 1-6, have been identified. Prefoldin 1 is a 122 amino acid protein that binds specifically to cytosolic chaperonin (c-cpn) and transfers target proteins to it. Prefoldin 3 (VBP1 or VHL binding protein-1) forms complexes with VHL and is translocated from perinuclear granules to the nucleus or cytoplasm. Prefoldin 4 is a possible transcription factor and prefoldin 5 is a c-Myc binding protein.

REFERENCES

1. Tsuchiya, H., Iseda, T. and Hino, O. 1996. Identification of a novel protein (VBP-1) binding to the von Hippel-Lindau (VHL) tumor suppressor gene product. *Cancer Res.* 56: 2881-2885.
2. Brinke, A., Green, P.M. and Giannelli, F. 1997. Characterization of the gene (VBP1) and transcript for the von Hippel-Lindau binding protein and isolation of the highly conserved murine homologue. *Genomics* 45: 105-112.
3. Vainberg, I.E., Lewis, S.A., Rommelaere, H., Ampe, C., Vandekerckhove, J., Klein, H.L. and Cowan, N.J. 1998. Prefoldin, a chaperone that delivers unfolded proteins to cytosolic chaperonin. *Cell* 93: 863-873.
4. LocusLink Report (LocusID: 300133). <http://www.ncbi.nlm.nih.gov/LocusLink/>
5. International Radiation Hybrid Mapping Consortium. (GenemapID:SHGC-31943) <http://www.ncbi.nlm.nih.gov/genemap/>

CHROMOSOMAL LOCATION

Genetic locus: PFDN6 (human) mapping to 6p21.32.

SOURCE

prefoldin 6 (1D10-4D11) is a mouse monoclonal antibody raised against prefoldin 6 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

prefoldin 6 (1D10-4D11) is available conjugated to either phycoerythrin (sc-53519 PE) or fluorescein (sc-53519 FITC), 200 µg/ml, for IF, IHC(P) and FCM.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

prefoldin 6 (1D10-4D11) is recommended for detection of prefoldin 6 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10⁶ cells).

Suitable for use as control antibody for prefoldin 6 siRNA (h): sc-40878, prefoldin 6 shRNA Plasmid (h): sc-40878-SH and prefoldin 6 shRNA (h) Lentiviral Particles: sc-40878-V.

Molecular Weight of prefoldin 6: 13 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RESEARCH USE

For research use only, not for use in diagnostic procedures.