



VCAM-1 (MR106): sc-53548

BACKGROUND

Cell adhesion molecules are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play an important role in embryogenesis and development. Neuronal cell adhesion molecule (NCAM) expression is observed in a variety of human tumors including neuroblastomas, rhabdomyosarcomas, Wilm's tumors, Ewing's sarcomas and some primitive myeloid malignancies. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the B2 α 1 and B2 α M integrins. PECAM-1 (platelet/endothelial cell adhesion molecule-1), also referred to as CD31, is a glycoprotein expressed on the cell surfaces of monocytes, neutrophils, platelets and a subpopulation of T cells. VCAM-1 (vascular cell adhesion molecule-1) was first identified as an adhesion molecule induced on human endothelial cells by inflammatory cytokines such as IL-1, tumor necrosis factor (TNF) and lipopolysaccharide (LPS). The KALIG gene encodes a nerve cell adhesion molecule (NCAM)-like protein and is deleted in 66% of patients with Kallmann's syndrome, anosmia with secondary hypogonadism.

REFERENCES

1. Thornhill, M.H., et al. 1991. Tumor necrosis factor combines with IL-4 or IFN- γ to selectively enhance endothelial cell adhesiveness for T cells. The contribution of vascular cell adhesion molecule-1-dependent and -independent binding mechanisms. *J. Immunol.* 146: 592-598.
2. Bevilacqua, M.P. 1993. Endothelial-leukocyte adhesion molecules. *Annu. Rev. Immunol.* 11: 767-804.
3. Cowen, M.A. and Green, M. 1993. The Kallmann's syndrome variant (KSV) model of the schizophrenias. *Schizophr. Res.* 9: 1-10.
4. Patel, K., et al. 1993. Vase mini-exon usage by NCAM is not restricted to tumours of neuroectodermal origin. *Int. J. Cancer* 54: 772-777.
5. Jorgensen, O.S. 1995. Neural cell adhesion molecule (NCAM) as a quantitative marker in synaptic remodeling. *Neurochem. Res.* 20: 533-547.
6. Edelman, G.M. and Jones, F.S. 1995. Developmental control of N-CAM expression by Hox and Pax gene products. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 349: 305-312.
7. Dominici, C., et al. 1996. Bone marrow micrometastases in a patient with localized Wilms' tumor. *Med. Pediatr. Oncol.* 26: 125-128.
8. Briskin, M.J., et al. 1996. Structural requirements for mucosal vascular addressin binding to its lymphocyte receptor α 4/ β 7. Common themes among integrin-Ig family interactions. *J. Immunol.* 156: 719-726.
9. Mayet, W.J., et al. 1996. Antibodies to proteinase 3 mediate expression of vascular cell adhesion molecule-1 (VCAM-1). *Clin. Exp. Immunol.* 103: 259-267.

CHROMOSOMAL LOCATION

Genetic locus: Vcam1 (mouse) mapping to 3 G1.

SOURCE

VCAM-1 (MR106) is a mouse monoclonal antibody raised against rat VCAM-1 from transfected mouse L5178Y cells.

PRODUCT

Each vial contains 200 μ g IgG κ light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

VCAM-1 (MR106) is available conjugated to either phycoerythrin (sc-53548 PE) or fluorescein (sc-53548 FITC), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM.

APPLICATIONS

VCAM-1 (MR106) is recommended for detection of VCAM-1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for VCAM-1 siRNA (m): sc-36810, VCAM-1 shRNA Plasmid (m): sc-36810-SH and VCAM-1 shRNA (m) Lentiviral Particles: sc-36810-V.

Molecular Weight of VCAM-1: 110 kDa.

Positive Controls: L8 cell lysate: sc-3807.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

SELECT PRODUCT CITATIONS

1. Taki-Eldin, A., et al. 2012. Triiodothyronine attenuates hepatic ischemia/reperfusion injury in a partial hepatectomy model through inhibition of proinflammatory cytokines, transcription factors, and adhesion molecules. *J. Surg. Res.* 178: 646-656.
2. Martiáñez, T., et al. 2013. N-cadherin expression is regulated by UTP in schwannoma cells. *Purinergic Signal.* 9: 259-270.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.