

LAT (11B.12): sc-53550



The Power to Question

BACKGROUND

T cell receptors activate immune responses by recognizing antigen and initiating a cascade of intracellular signal transduction events, eventually culminating in cell proliferation and differentiation. Both protein tyrosine kinases and PLC γ are activated by this event. LAT, or linker for activation of T cells, is an integral membrane protein that has been shown to associate with PLC γ 1, as well as GRB2 and the p85 subunit of PI 3-kinase. Binding of these signaling molecules to LAT is associated with phosphorylation of LAT by ZAP-70/Syk tyrosine kinases. LAT appears to play a role in activation of transcription mediated by AP-1 and NFAT following stimulation of the T cell receptor, suggesting that it acts as a linker protein in T cell activation. LAT protein is palmitoylated, and this modification is required for its tyrosine phosphorylation and localization to glycolipid-enriched microdomains.

CHROMOSOMAL LOCATION

Genetic locus: LAT (human) mapping to 16p11.2; Lat (mouse) mapping to 7 F3.

SOURCE

LAT (11B.12) is a mouse monoclonal antibody raised against amino acids 31-233 of LAT of human origin.

PRODUCT

Each vial contains 200 μ g IgG κ light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

LAT (11B.12) is available conjugated to agarose (sc-53550 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-53550 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-53550 PE), fluorescein (sc-53550 FITC), Alexa Fluor[®] 488 (sc-53550 AF488), Alexa Fluor[®] 546 (sc-53550 AF546), Alexa Fluor[®] 594 (sc-53550 AF594) or Alexa Fluor[®] 647 (sc-53550 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-53550 AF680) or Alexa Fluor[®] 790 (sc-53550 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

LAT (11B.12) is recommended for detection of LAT of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for LAT siRNA (h): sc-35795, LAT siRNA (m): sc-35796, LAT shRNA Plasmid (h): sc-35795-SH, LAT shRNA Plasmid (m): sc-35796-SH, LAT shRNA (h) Lentiviral Particles: sc-35795-V and LAT shRNA (m) Lentiviral Particles: sc-35796-V.


Molecular Weight of LAT: 36-38 kDa.

Positive Controls: WR19L cell lysate: sc-3805, BYDP whole cell lysate: sc-364368 or Jurkat whole cell lysate: sc-2204.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



LAT (11B.12) Alexa Fluor[®] 647: sc-53550 AF647. Direct fluorescent western blot analysis of LAT expression in BYDP (A), WR19L (B), Jurkat (C) and ALL-SIL (D) whole cell lysates. Blocked with UltraCruz[®] Blocking Reagent: sc-516214. Cruz Marker[™] Molecular Weight Standards detected with Cruz Marker[™] MW Tag-Alexa Fluor[®] 488: sc-516790.



LAT (11B.12): sc-53550. Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing membrane and cytoplasmic staining of cells in white pulp and cells in red pulp.

SELECT PRODUCT CITATIONS

- Mingueneau, M., et al. 2009. Loss of the LAT adaptor converts antigen-responsive T cells into pathogenic effectors that function independently of the T cell receptor. *Immunity* 31: 197-208.
- Tai, T.S., et al. 2013. GATA-3 regulates the homeostasis and activation of CD8⁺ T cells. *J. Immunol.* 190: 428-437.
- Yang, Y., et al. 2015. The Us3 protein of herpes simplex virus 1 inhibits T cell signaling by confining linker for activation of T cells (LAT) activation via TRAF6 protein. *J. Biol. Chem.* 290: 15670-15678.
- Nguyen, N.N.T., et al. 2018. Hepatitis C virus modulates solute carrier family 3 member 2 for viral propagation. *Sci. Rep.* 8: 15486.
- Yang, T., et al. 2020. Transcriptomic signature of gut microbiome-contacting cells in colon of spontaneously hypertensive rats. *Physiol. Genomics* 52: 121-132.
- Sugimoto, C., et al. 2022. Reprogramming and redifferentiation of mucosal-associated invariant T cells reveal tumor inhibitory activity. *Elife* 11: e70848.
- Kim, H.W., et al. 2022. NAD⁺-boosting molecules suppress mast cell degranulation and anaphylactic responses in mice. *Theranostics* 12: 3316-3328.
- Saitoh, K., et al. 2022. STAP-2 is a novel positive regulator of TCR-proximal signals. *J. Immunol.* 209: 57-68.
- Chang, H.W., et al. 2023. Thalidomide attenuates mast cell activation by upregulating SHP-1 signaling and interfering with the action of CRBN. *Cells* 12: 469.
- Yang, X., et al. 2024. Cannabidiol inhibits IgE-mediated mast cell degranulation and anaphylaxis in mice. *Mol. Nutr. Food Res.* 68: e2300136.

RESEARCH USE

For research use only, not for use in diagnostic procedures.