

CD7 (SBCV9.1): sc-53568

BACKGROUND

CD7 (also designated T cell leukemia antigen) is a type I transmembrane glycoprotein that is expressed on pluripotent hemopoietic cells, most human thymocytes and some peripheral blood T cells. CD7 is a marker for pluripotent stem cell leukemias and T cell acute lymphocytic leukemia. A role for CD7 in the activation of T cells with γ/δ receptors has been suggested. CD8 T cells from patients infected with HIV-1 displayed profound down-modulation of CD7 expression as compared with healthy subjects. CD7 is among the pan-T-cell antigens down-regulated in acute infectious mononucleosis.

REFERENCES

- Haynes, B.F., et al. 1989. Ontogeny of T cell precursors: a model for the initial stages of human T cell development. *Immunol. Today* 10: 87-91.
- Barcena, A., et al. 1995. Tracing the expression of CD7 and other antigens during T and myeloid cell differentiation in the human fetal liver and thymus. *Leuk. Lymphoma* 17: 1-11.
- Schanberg, L.E., et al. 1995. Characterization of human CD7 transgenic mice. *J. Immunol.* 155: 2407-2418.
- Leta, E., et al. 1995. Production and characterization of the extracellular domain of human CD7 antigen: further evidence that CD7 has a role in T cell signaling. *Cell. Immunol.* 165: 101-109.
- Ward, S.G., et al. 1995. Antibody ligation of CD7 leads to association with phosphoinositide 3-kinase and phosphatidylinositol 3,4,5-triphosphate formation in T lymphocytes. *Eur. J. Immunol.* 25: 502-507.
- Weisberger, J., et al. 2003. Down-regulation of pan-T-cell antigens, particularly CD7, in acute infectious mononucleosis. *Am. J. Clin. Pathol.* 120: 49-55.
- Tiftik, N., et al. 2004. The importance of CD7 and CD56 antigens in acute leukaemias. *Int. J. Clin. Pract.* 58: 149-152.
- Aandahl, E.M., et al. 2004. Expansion of CD7(low) and CD7(negative) CD8 T cell effector subsets in HIV-1 infection: correlation with antigenic load and reversion by antiretroviral treatment. *Blood* 104: 3672-3678.

CHROMOSOMAL LOCATION

Genetic locus: CD7 (human) mapping to 17q25.3.

SOURCE

CD7 (SBCV9.1) is a mouse monoclonal antibody raised against peripheral blood mononuclear leukocytes of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CD7 (SBCV9.1) is recommended for detection of CD7 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for CD7 siRNA (h): sc-35021, CD7 shRNA Plasmid (h): sc-35021-SH and CD7 shRNA (h) Lentiviral Particles: sc-35021-V.


Molecular Weight of CD7: 40 kDa.

Positive Controls: CCRF-CEM cell lysate: sc-2225, Jurkat whole cell lysate: sc-2204 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CD7 (SBCV9.1): sc-53568. Indirect FCM analysis of human peripheral blood leukocytes stained with CD7 (SBCV9.1), followed by PE-conjugated goat anti-mouse IgG: sc-3738. Black line histogram represents the isotype control, normal mouse IgG: sc-3879.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.