

NKp44 (3.43): sc-53598

BACKGROUND

Natural killer (NK) cells direct cytotoxicity against tumor or virally infected cells. NK cell-mediated cytotoxicity is stimulated by several activating receptors associated with the signaling adapter DNAX activation 12/killer cell-activating receptor-associated protein (DAP12). NKp44 is a natural cytotoxicity receptor that is expressed on IL-2-activated human NK cells and may contribute to the increased efficiency of NK cells to mediate tumor cell lysis. NKp44 is composed of one Ig-like extracellular domain, a transmembrane segment and a cytoplasmic domain. Prolactin upregulates and cortisol downregulates the surface expression of NKp44 at the transcriptional level. A cellular ligand for NKp44 (NKp44L) is expressed during HIV-1 infection and is correlated with the progression of CD4⁺ T cell depletion and an increase of viral load. This implicates NKp44 as a therapeutic agent that may aid in the progress towards a vaccine for HIV-1 infection.

REFERENCES

- Cantoni, C., Ponassi, M., Biassoni, R., Conte, R., Spallarossa, A., Moretta, A., Moretta, L., Bolognesi, M. and Bordo, D. 2003. The three-dimensional structure of the human NK cell receptor NKp44, a triggering partner in natural cytotoxicity. *Structure* 11: 725-734.
- De Maria, A., Fogli, M., Costa, P., Murdaca, G., Puppo, F., Mavilio, D., Moretta, A. and Moretta, L. 2003. The impaired NK cell cytolytic function in viremic HIV-1 infection is associated with a reduced surface expression of natural cytotoxicity receptors (NKp46, NKp30 and NKp44). *Eur. J. Immunol.* 33: 2410-2418.
- Campbell, K.S., Yusa, S., Kikuchi-Maki, A. and Catina, T.L. 2004. NKp44 triggers NK cell activation through DAP12 association that is not influenced by a putative cytoplasmic inhibitory sequence. *J. Immunol.* 172: 899-906.
- Forte, P., Lilienfeld, B.G., Baumann, B.C. and Seebach, J.D. 2005. Human NK cytotoxicity against porcine cells is triggered by NKp44 and NKG2D. *J. Immunol.* 175: 5463-5470.
- Fuchs, A., Cella, M., Kondo, T. and Colonna, M. 2005. Paradoxical inhibition of human natural interferon-producing cells by the activating receptor NKp44. *Blood* 106: 2076-2082.
- Mavoungou, E., Bouyou-Akotet, M.K. and Kreamsner, P.G. 2005. Effects of prolactin and cortisol on natural killer (NK) cell surface expression and function of human natural cytotoxicity receptors (NKp46, NKp44 and NKp30). *Clin. Exp. Immunol.* 139: 287-296.
- Srivastava, B.I. and Srivastava, M.D. 2005. Expression of natural cytotoxicity receptors NKp30, NKp44, and NKp46 mRNAs and proteins by human hematopoietic and non-hematopoietic cells. *Leuk. Res.* 30: 37-46.
- Stet, R.J., Hermsen, T., Westphal, A.H., Jukes, J., Engelsma, M., Lidy Verburg-van Kemenade, B.M., Dortmans, J., Aveiro, J. and Savelkoul, H.F. 2005. Novel immunoglobulin-like transcripts in teleost fish encode polymorphic receptors with cytoplasmic ITAM or ITIM and a new structural Ig domain similar to the natural cytotoxicity receptor NKp44. *Immunogenetics* 57: 77-89.

CHROMOSOMAL LOCATION

Genetic locus: NCR2 (human) mapping to 6p21.1.

SOURCE

NKp44 (3.43) is a mouse monoclonal antibody raised against NKp44 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NKp44 (3.43) is available conjugated to either phycoerythrin (sc-53598 PE) or fluorescein (sc-53598 FITC), 200 µg/ml, for IF, IHC(P) and FCM.

APPLICATIONS

NKp44 (3.43) is recommended for detection of NKp44 of human origin by flow cytometry (1 µg per 1 x 10⁶ cells).

Suitable for use as control antibody for NKp44 siRNA (h): sc-72170, NKp44 shRNA Plasmid (h): sc-72170-SH and NKp44 shRNA (h) Lentiviral Particles: sc-72170-V.

Molecular Weight of NKp44: 44 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.