

IgG_{2c} (MRG2c-67): sc-53742

BACKGROUND

IgG is a monomeric immunoglobulin composed of two heavy chains and two light chains. There are 4 subclasses of IgG: IgG₁, IgG₂, IgG₃ and IgG₄. Each molecule has two antigen binding sites. IgG is the most abundant immunoglobulin as well as the only isotype that can pass through the placenta, thereby providing protection to the fetus in its first weeks of life before its own immune system has developed. IgG can bind to several different kinds of pathogens, for example viruses, bacteria and fungi and it protects the body against them by complement activation (the classic pathway), opsonization for phagocytosis and neutralization of their toxins.

REFERENCES

1. Adetugbo, K. 1978. Evolution of immunoglobulin subclasses. Primary structure of a murine myeloma γ 1 chain. *J. Biol. Chem.* 253: 6068-6075.
2. Tucker, P.W., Marcu, K.B., Slightom, J.L. and Blattner, F.R. 1979. Structure of the constant and 3' untranslated regions of the murine γ 2b heavy chain messenger RNA. *Science* 206: 1299-1303.
3. Rabbitts, T.H., Forster, A., Dunnick, W. and Bentley, D.L. 1980. The role of gene deletion in the immunoglobulin heavy chain switch. *Nature* 283: 351-356.
4. Sakano, H., Maki, R., Kurosawa, Y., Roeder, W. and Tonegawa, S. 1980. Two types of somatic recombination are necessary for the generation of complete immunoglobulin heavy-chain genes. *Nature* 286: 676-683.
5. Goldsby, R., Kindt, T. and Osborne, B. 1992. *Immunology*. New York: W.H. Freeman and Company.
6. Wuhrer, M., Stam, J.C., van de Geijn, F.E., Koeleman, C.A., Verrips, C.T., Dolhain, R.J., Hokke, C.H. and Deelder, A.M. 2007. Glycosylation profiling of immunoglobulin G (IgG) subclasses from human serum. *Proteomics* 7: 4070-4081.
7. Nair, N., Gans, H., Lew-Yasukawa, L., Long-Wagar, A.C., Arvin, A. and Griffin, D.E. 2007. Age-dependent differences in IgG isotype and avidity induced by measles vaccine received during the first year of life. *J. Infect. Dis.* 196: 1339-1345.
8. Fuchs, S., Feferman, T., Zhu, K.Y., Meidler, R., Margalit, R., Wang, N., Laub, O. and Souroujon, M.C. 2007. Suppression of experimental autoimmune myasthenia gravis by intravenous immunoglobulin and isolation of a disease-specific IgG fraction. *Ann. N.Y. Acad. Sci.* 1110: 550-558.

SOURCE

IgG_{2c} (MRG2c-67) is a mouse monoclonal antibody raised against mixed immunoglobulins of rat origin.

PRODUCT

Each vial contains 100 μ g IgG_{2c} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

IgG_{2c} (MRG2c-67) is recommended for detection of IgG_{2c} of mouse and rat origin by flow cytometry (1 μ g per 1 x 10⁶ cells); non cross-reactive with other isotypes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.