

CD2 (1E7E8): sc-53823

BACKGROUND

CD2 (also designated E-rosette receptor) interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes, as well as on mouse B cells and rat splenic macrophages. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counterreceptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

REFERENCES

1. Shaw, A.S., et al. 1997. Making the T cell receptor go the distance: a topological view of T cell activation. *Immunity* 6: 361-369.
2. Dustin, M.L., et al. 1998. A novel adaptor protein orchestrates receptor patterning and cytoskeletal polarity in T cell contacts. *Cell* 94: 667-677.
3. Nishizawa, K., et al. 1998. Identification of a proline-binding motif regulating CD2-triggered T lymphocyte activation. *Proc. Natl. Acad. Sci. USA* 95: 14897-14902.
4. Shih, N.Y., et al. 1999. Congenital nephrotic syndrome in mice lacking CD2-associated protein. *Science* 286: 312-315.
5. Guan, F., et al. 2006. Autocrine VEGF-A system in podocytes regulates podocin and its interaction with CD2AP. *Am. J. Physiol. Renal Physiol.* 291: F422-F428.
6. Fan, Q., et al. 2006. The relationship among nephrin, podocin, CD2AP and α -actinin might not be a true "interaction" in podocyte. *Kidney Int.* 69: 1207-1215.
7. Xia, W., et al. 2006. Differential interactions between transforming growth factor β 3/ β R1, TAB1 and CD2AP disrupt blood-testis barrier and Sertoli-germ cell adhesion. *J. Biol. Chem.* 281: 16799-16813.

CHROMOSOMAL LOCATION

Genetic locus: CD2 (human) mapping to 1p13.1.

SOURCE

CD2 (1E7E8) is a mouse monoclonal antibody raised against thymocytes/Sezary T cells of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD2 (1E7E8) is available conjugated to either phycoerythrin (sc-53823 PE) or fluorescein (sc-53823 FITC), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CD2 (1E7E8) is recommended for detection of CD2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for CD2 siRNA (h): sc-29970, CD2 shRNA Plasmid (h): sc-29970-SH and CD2 shRNA (h) Lentiviral Particles: sc-29970-V.

Molecular Weight of CD2: 50 kDa.


Positive Controls: CD2 (h2): 293T Lysate: sc-172563, HuT 78 whole cell lysate: sc-2208 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



CD2 (1E7E8): sc-53823. Western blot analysis of CD2 expression in non-transfected 293T: sc-117752 (A), human CD2 transfected 293T: sc-172563 (B) and Jurkat (C) whole cell lysates.



CD2 (1E7E8): sc-53823. Indirect FCM analysis of human peripheral blood leukocytes stained with CD2 (1E7E8), followed by PE-conjugated goat anti-mouse IgG: sc-3738. Black line histogram represents the isotype control, normal mouse IgG_{2a}: sc-3878.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.