EMMPRIN (RL73): sc-53862



The Power to Question

BACKGROUND

Extracellular matrix metalloproteinase inducer, EMMPRIN (also designated basigin or CD147), is involved in the regulation of matrix remodeling at the epidermal-dermal interface. EMMPRIN stimulates the production of interstitial collagenase, gelatinase A, stromelysin-1 and various metalloproteinases (MMPs) by fibroblasts. These enzymes, which are typically increased during tissue degradation and wound healing, are important factors in cancer invasion and metastasis.

REFERENCES

- 1. Paterson, D.J., et al. 1987. Antigens of activated rat T lymphocytes including a molecule of 50,000 $\rm M_r$ detected only on CD4 positive T blasts. Mol. Immunol. 24: 1281-1290.
- Miyauchi, T., et al. 1990. Basigin, a new, broadly distributed member of the immunoglobulin superfamily, has strong homology with both the immunoglobulin V domain and the β chain of major histocompatibility complex class II antigen. J. Biochem. 107: 316-323.
- Biswas, C., et al. 1995. The human tumor cell-derived collagenase stimulatory factor (renamed EMMPRIN) is a member of the immunoglobulin superfamily. Cancer Res. 55: 434-439.
- DeCastro, R., et al. 1996. Human keratinocytes express EMMPRIN, an extracellular matrix metalloproteinase inducer. J. Invest. Dermatol. 106: 1260-1265.
- Guo, H., et al. 1997. Stimulation of matrix metalloproteinase production by recombinant extracellular matrix metalloproteinase inducer from transfected Chinese hamster ovary cells. J. Biol. Chem. 272: 24-27.
- Guo, H., et al. 1998. Characterization of the gene for human EMMPRIN, a tumor cell surface inducer of matrix metalloproteinases. Gene 220: 99-108.
- Lim, M., et al. 1998. Tumor-derived EMMPRIN (extracellular matrix metalloproteinase inducer) stimulates collagenase transcription through MAPK p38. FEBS Lett. 441: 88-92.
- 8. Kahari, V.M., et al. 1999. Matrix metalloproteinases and their inhibitors in tumour growth and invasion. Ann. Med. 31: 34-45.

CHROMOSOMAL LOCATION

Genetic locus: Bsg (mouse) mapping to 10 C1.

SOURCE

EMMPRIN (RL73) is a rat monoclonal antibody raised against EL4-6.1 cell line of mouse origin.

PRODUCT

Each vial contains 200 μg lgG_{2a} in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

EMMPRIN (RL73) is available conjugated to either phycoerythrin (sc-53862 PE) or fluorescein (sc-53862 FITC), 200 μg/ml, for WB (RGB), IF, IHC(P) and FCM.

APPLICATIONS


EMMPRIN (RL73) is recommended for detection of EMMPRIN of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).


Suitable for use as control antibody for EMMPRIN siRNA (m): sc-35299, EMMPRIN shRNA Plasmid (m): sc-35299-SH and EMMPRIN shRNA (m) Lentiviral Particles: sc-35299-V.


Molecular Weight of EMMPRIN: 55 kDa.

Positive Controls: RAW 264.7 whole cell lysate: sc-2211, CTLL-2 cell lysate: sc-2242 or MH-S whole cell lysate: sc-364785.

DATA







EMMPRIN (RL73): sc-53862. Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse placenta tissue showing membrane and cytoplasmic staining of trophoblastic cells.

SELECT PRODUCT CITATIONS

 Ozeki, N., et al. 2015. A2 integrin, extracellular matrix metalloproteinase inducer, and matrix metalloproteinase-3 act sequentially to induce differentiation of mouse embryonic stem cells into odontoblast-like cells. Exp. Cell Res. 331: 21-37.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **EMMPRIN (B-5): sc-46700** for EMMPRIN antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor* 488, 546, 594, 647, 680 and 790.