

ZAP-70 (3D2A4): sc-53899

BACKGROUND

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR) which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the T α and β chains, the CD3 γ , δ and ϵ chains and a ζ -containing homodimer or heterodimer. The disulfide-linked T α - β heterodimer is responsible for antigen recognition, but the short 5 amino acid cytoplasmic domains of T α and β are unlikely to be sufficient to couple to intracellular signaling pathways. In contrast, the structured features of the CD3 and ζ subunits suggest a role in signal transduction. Of these, the ζ chain, which is expressed as either a homodimer or heterodimer, has a short extracellular domain of only 9 amino acids, but a larger 113 amino acid cytoplasmic domain. A tyrosine phosphoprotein, ZAP-70, has been identified that associates with ζ and undergoes tyrosine phosphorylation following TCR stimulation.

REFERENCES

1. Clevers, H., et al. 1988. The T cell receptor/CD3 complex: a dynamic protein ensemble. *Annu. Rev. Immunol.* 6: 629-662.
2. Baniyash, M., et al. 1988. The T cell antigen receptor ζ chain is tyrosine phosphorylated upon activation. *J. Biol. Chem.* 263: 18225-18230.
3. Baniyash, M., et al. 1989. The isolation and characterization of the murine T cell antigen receptor ζ chain gene. *J. Biol. Chem.* 264: 13252-13257.
4. Frank, S.J., et al. 1990. The structure and signaling function of the invariant T cell receptor components. *Semin. Immunol.* 2: 89-97.
5. Clayton, L.K., et al. 1991. CD3 η and CD3 ζ are alternatively spliced products of a common genetic locus and are transcriptionally and/or post-transcriptionally regulated during T-cell development. *Proc. Natl. Acad. Sci. USA* 88: 5202-5206.
6. Chan, A.C., et al. 1991. The ζ chain is associated with a tyrosine kinase and upon T-cell antigen receptor stimulation associates with ZAP-70, a 70-kDa tyrosine phosphoprotein. *Proc. Natl. Acad. Sci. USA* 88: 9166-9170.
7. Chan, A.C., et al. 1992. ZAP-70: a 70 kd protein-tyrosine kinase that associates with the TCR ζ chain. *Cell* 71: 649-662.

CHROMOSOMAL LOCATION

Genetic locus: ZAP70 (human) mapping to 2q11.2; Zap70 (mouse) mapping to 1 B.

SOURCE

ZAP-70 (3D2A4) is a mouse monoclonal antibody raised against truncated recombinant ZAP-70 of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_{2a}$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ZAP-70 (3D2A4) is recommended for detection of ZAP-70 of mouse, rat and human origin by Western Blotting (starting dilution 1:500, dilution range 1:500-1:2,000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for ZAP-70 siRNA (h): sc-29526, ZAP-70 siRNA (m): sc-36867, ZAP-70 shRNA Plasmid (h): sc-29526-SH, ZAP-70 shRNA Plasmid (m): sc-36867-SH, ZAP-70 shRNA (h) Lentiviral Particles: sc-29526-V and ZAP-70 shRNA (m) Lentiviral Particles: sc-36867-V.

Molecular Weight of ZAP-70: 70 kDa.


Positive Controls: ZAP-70 (h): 293T Lysate: sc-114635, Jurkat whole cell lysate: sc-2204 or MOLT-4 cell lysate: sc-2233.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ZAP-70 (3D2A4): sc-53899. Western blot analysis of ZAP-70 expression in non-transfected 293T: sc-117752 (A), human ZAP-70 isoform 3 transfected 293T: sc-114635 (B) and Jurkat (C) whole cell lysates.



ZAP-70 (3D2A4): sc-53899. Western blot analysis of ZAP-70 expression in MOLT-4 (A), BYDP (B) and CTL-2 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **ZAP-70 (1E7.2): sc-32760** for ZAP-70 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.