

Mcl-1 (8C6D4B1): sc-53951

BACKGROUND

B cell CLL/lymphoma-2 (Bcl-2) blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal. Myeloid cell leukemia-1 (Mcl-1) shares sequence homology with Bcl-2 and further resembles Bcl-2 in that its expression promotes cell viability. P53 and Mcl-1 demonstrate opposing effects on mitochondrial apoptosis by mediating Bcl-2 antagonist killer (Bak) activity. Mcl-1 is an important and specific regulator that is necessary for the homeostasis of early hematopoietic progenitors. Glycogen synthase kinase-3 (GSK-3) controls Mcl-1 stability, which has an effect on the regulation of apoptosis by growth factors PI 3-kinase and Akt. Mice with a deficiency of the Mcl-1 protein show a significant reduction in B and T lymphocytes similar to the effects observed in IL-7- or IL-7R-deficient mice.

CHROMOSOMAL LOCATION

Genetic locus: MCL1 (human) mapping to 1q21.3; Mcl1 (mouse) mapping to 3 F2.1.

SOURCE

Mcl-1 (8C6D4B1) is a mouse monoclonal antibody raised against recombinant Mcl-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Mcl-1 (8C6D4B1) is recommended for detection of Mcl-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Mcl-1 siRNA (h): sc-35877, Mcl-1 siRNA (m): sc-35878, Mcl-1 shRNA Plasmid (h): sc-35877-SH, Mcl-1 shRNA Plasmid (m): sc-35878-SH, Mcl-1 shRNA (h) Lentiviral Particles: sc-35877-V and Mcl-1 shRNA (m) Lentiviral Particles: sc-35878-V.

Molecular Weight of Mcl-1 long form: 40 kDa.


Molecular Weight of Mcl-1 short form: 32 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HL-60 whole cell lysate: sc-2209 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Mcl-1 (8C6D4B1): sc-53951. Western blot analysis of Mcl-1 expression in F9 whole cell lysate.



Mcl-1 (8C6D4B1): sc-53951. Immunoperoxidase staining of formalin fixed, paraffin-embedded human appendix tissue showing cytoplasmic staining of glandular cells and lymphoid cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing cytoplasmic staining of trophoblastic cells (B).

SELECT PRODUCT CITATIONS

- Kim, J.H. and Park, B. 2017. Triptolide blocks the Stat3 signaling pathway through induction of protein tyrosine phosphatase SHP-1 in multiple myeloma cells. *Int. J. Mol. Med.* 40: 1566-1572.
- Luo, G., et al. 2018. Dendritic cell factor 1 inhibits proliferation and migration and induces apoptosis of neuroblastoma cells by inhibiting the ERK signaling pathway. *Oncol. Rep.* 41: 103-112.
- Wudtiwai, B., et al. 2018. α-mangostin, an active compound in *Garcinia mangostana*, abrogates anoikis-resistance in human hepatocellular carcinoma cells. *Toxicol. In Vitro* 53: 222-232.
- Ma, Z. and Xue, X. 2018. Differentially expressed proteins in the human esophageal cancer cell line Eca-109, in the presence and absence of gemcitabine. *Mol. Med. Rep.* 17: 1873-1878.
- Liu, S.J., et al. 2018. MiR-15a-3p affects the proliferation, migration and apoptosis of lens epithelial cells. *Mol. Med. Rep.* 19: 1110-1116.



See **Mcl-1 (22): sc-12756** for Mcl-1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.