

Mcl-1 (RC31): sc-53971

BACKGROUND

B-cell CLL/lymphoma 2 (Bcl-2) blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal. Myeloid cell leukemia 1 (Mcl-1) shares sequence homology with Bcl-2 and further resembles Bcl-2 in that its expression promotes cell viability. p53 and Mcl-1 demonstrate opposing effects on mitochondrial apoptosis by mediating Bcl-2 antagonist killer (Bak) activity. Mcl-1 is an important and specific regulator that is necessary for the homeostasis of early hematopoietic progenitors. Glycogen synthase kinase 3 (GSK3) controls Mcl-1 stability, which has an effect on the regulation of apoptosis by growth factors, PI 3-kinase and Akt. Mice with a deficiency of the Mcl-1 protein show a significant reduction in B and T lymphocytes similar to the effects observed in IL-7- or IL-7R-deficient mice.

REFERENCES

1. Kozopas, K.M., et al. 1993. Mcl-1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to Bcl-2. *Proc. Natl. Acad. Sci. USA* 90: 3516-3520.
2. Craig, R.W., et al. 1994. Human and mouse chromosomal mapping of the myeloid cell leukemia-1 gene: Mcl-1 maps to human chromosome 1q21, a region that is frequently altered in preneoplastic and neoplastic disease. *Genomics* 23: 457-463.
3. Rinkenberger, J.L., et al. 2000. Mcl-1 deficiency results in peri-implantation embryonic lethality. *Genes Dev.* 14: 23-27.
4. Bae, J., et al. 2000. Mcl-1S, a splicing variant of the antiapoptotic Bcl-2 family member Mcl-1, encodes a proapoptotic protein possessing only the BH3 domain. *J. Biol. Chem.* 275: 25255-25261.
5. Opferman, J.T., et al. 2003. Development and maintenance of B and T lymphocytes requires antiapoptotic Mcl-1. *Nature* 426: 671-676.
6. Leu, J.L., et al. 2004. Mitochondrial p53 activates Bak and causes disruption of a Bak-Mcl-1 complex. *Nat. Cell Biol.* 6: 443-450.
7. Opferman, J.T., et al. 2005. Obligate role of anti-apoptotic Mcl-1 in the survival of hematopoietic stem cells. *Science* 307: 1101-1104.

CHROMOSOMAL LOCATION

Genetic locus: MCL1 (human) mapping to 1q21.3.

SOURCE

Mcl-1 (RC31) is a mouse monoclonal antibody raised against amino acids 1-327 of Mcl-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Mcl-1 (RC31) is recommended for detection of Mcl-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Mcl-1 siRNA (h): sc-35877, Mcl-1 shRNA Plasmid (h): sc-35877-SH and Mcl-1 shRNA (h) Lentiviral Particles: sc-35877-V.

Molecular Weight of Mcl-1 long form: 40 kDa.


Molecular Weight of Mcl-1 short form: 32 kDa.

Positive Controls: BJAB whole cell lysate: sc-2207, Ramos cell lysate: sc-2216 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Mcl-1 (RC31): sc-53971. Western blot analysis of Mcl-1 expression in BJAB (A), Ramos (B) and K-562 (C) whole cell lysates.



Mcl-1 (RC31): sc-53971. Western blot analysis of Mcl-1 expression in HeLa (A), Raji (B) and Ramos (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Mcl-1 (22): sc-12756** for Mcl-1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.